

CURRENT RESEARCH, MONITORING, AND EDUCATION PROJECTS

2018 – 2019

Baruch Marine Field Laboratory

**North Inlet-Winyah Bay
National Estuarine Research Reserve**

University of South Carolina



**Belle W. Baruch Institute
for Marine & Coastal Sciences**



**North Inlet-Winyah Bay
National Estuarine Research Reserve**

Current Research, Monitoring, and Education Projects 2018 – 2019

Introduction

The Baruch Marine Field Laboratory (BMFL), located on Hobcaw Barony in Georgetown County, has been the center of research activities for scientists and students from the University of South Carolina (USC) and dozens of other institutions since 1969. We conservatively estimate that between senior scientist projects and masters and doctoral studies conducted by graduate students, more than 1,000 grant and institutionally-funded projects have taken place at BMFL. This work has contributed substantially to the more than 2,000 peer-reviewed scientific articles, books, and technical reports that have been published since the Baruch Institute was founded. Independent and multi-disciplinary studies have been conducted by biologists, chemists, geologists, oceanographers, and other specialists who share interests in the structure, function, and condition of coastal environments. Results of research projects are used by educators, coastal resource managers, health and environmental regulators, legislators, and many other individuals and organizations interested in maintaining and improving the condition of estuaries in the face of increasing human activities and changing climate in the coastal zone.

The following annotated list summarizes 79 projects that were underway during the period from July 2018 through June 2019 in the North Inlet estuary and Winyah Bay by faculty, graduate students, staff, and undergraduates associated with the USC and other institutions. USC is the home institution for 53 of the investigators while over 70 investigators representing at least 30 other institutions and agencies are carrying out projects through BMFL. Dozens of graduate and undergraduate students assist scientists throughout the year to obtain hands-on training in field methods and to conduct research.

This annual report lists active projects (in random order) along with a project summary that includes the title, investigators, affiliations, and an abstract. Many of the studies that involve field measurements and collections are being conducted within the North Inlet–Winyah Bay National Estuarine Research Reserve (NI–WB NERR).

Funds for these research projects are provided by a variety of sources, including the National Science Foundation (NSF), Environmental Protection Agency (EPA), National Oceanic and Atmospheric Administration (NOAA) National Estuarine Research Reserve System (NERRS), South Carolina Sea Grant Consortium, US Department of Energy (US DOE), US Department of Defense (DoD), Office of Naval Research (ONR), National Aeronautics and Space Administration (NASA), Slocum-Lunz Foundation, Nemours Wildlife Foundation, South Carolina Independent Colleges and Universities, and the South Carolina Department of Health and Environmental Control (SC DHEC). The Belle W. Baruch Foundation provides the long-term stewardship of Hobcaw Barony, maintaining it in a natural state for research and education.

Contents

Map	9
Fluorescent dissolved organic matter dynamics in the North Inlet estuary	10
Investigators: Erik M. Smith, Julie L. Krask, Baker Stevens, Tracy Buck, and Susan Denham	
Ecology, behavior, and population biology of bottlenose dolphins (<i>Tursiops truncatus</i>) in the North Inlet estuary and adjacent waters	10
Investigators: Rob Young and students	
Interspecific competition among some salt marsh perennials in South Carolina	10
Investigators: Richard Stalter and John Baden	
Construction of biodegradable floating treatment wetland utilizing wood and bamboo	11
Investigators: William Strosnider, Travis Tasker, Lily Currie, and Kendra Zaruba	
A forty-seven year comparison of the vascular flora at three abandoned rice fields, Georgetown, South Carolina	11
Investigators: Richard Stalter, Joseph Rachlin, and John Baden	
Fish and crustacean use of marshes and intertidal creeks: Population and community level changes and relationships with weather and climate-driven changes	11
Investigators: Bruce W. Pfirrmann, Matthew E. Kimball, Dennis M. Allen, and Paul D. Kenny	
The conservation status of the canebrake rattlesnake (<i>Crotalus horridus atricaudatus</i>) at Hobcaw Barony, with identification of key areas for conservation of its herpetofauna	12
Investigator: Allan L. Markezich	
Seasonal energetics of the blue crab (<i>Callinectes sapidus</i>)	12
Investigators: Blaine D. Griffen, Matthew E. Kimball, and Bruce W. Pfirrmann	
Chemical characteristics of estuarine waters: Long-term monitoring at four sites in North Inlet estuary	12
Investigators: Erik M. Smith, Julie L. Krask, Baker Stevens, and Susan Denham	
Maintenance and operation of IOOS/SECOORA priority WERA HF radar sites	13
Investigators: George Voulgaris and William H. Jefferson	
Painted Bunting (<i>Passerina ciris</i>) monitoring project	13
Investigators: Jennifer Plunket and citizen science volunteers Wendy Allen, Bill Brabson, Cindy Kerr, Marlene Konsek, Pete Little, and Sandy Little	
Eddy covariance flux measurements to quantify salt marsh productivity and its response to environmental variability over multiple time scales	13
Investigators: Thomas L. O'Halloran and Erik M. Smith	
Benthic bivalves as potential indicator species for ecosystem climate change effects	14
Investigators: Juliana M. Harding and Dennis M. Allen	
Coastal training activities in the North Inlet–Winyah Bay National Estuarine Research Reserve: Protecting water and habitat quality through science-based community training	14
Investigator: Maeve Snyder	
Toxicological effects of tire wear particles on fathead minnow (<i>Pimephales promelas</i>) and Atlantic killifish (<i>Fundulus heteroclitus</i>)	14
Investigators: Peter van den Hurk, Stephanie LaPlaca, and John Weinstein	
Public and K-12 education activities: North Inlet–Winyah Bay National Estuarine Research Reserve	15
Investigators: Beth Thomas and Hayley Fournier	

Saltwater intrusion monitoring	15
Investigators: Alicia Wilson and William Clendenin	
Long-term changes in the zooplankton of the North Inlet estuary and relationships with climate change.....	15
Investigators: Dennis M. Allen, Josh Stone, Bruce W. Pfirrmann, and Matthew E. Kimball	
Quantifying elevational and latitudinal gradients in marsh snail predation.....	16
Investigators: Rebecca Atkins, Craig Osenberg, and Daniel Hawkins	
Mechanisms for thermal tolerance in an estuarine cnidarian	16
Investigator: Adam Reitzel	
Green porcelain crab (<i>Petrolisthes armatus</i>) larval biology and phenology	16
Investigator: Juliana M. Harding	
Exploring macroinvertebrate diversity of non-native macroalgae <i>Agarophyton vermiculophyllum</i>	16
Investigator: Timothy Lee	
Sediment accretion in North Inlet estuary salt marshes	17
Investigators: James T. Morris and Karen Sundberg	
Analyzing nickel, zinc, and lead in <i>Fundulus heteroclitus</i>.....	17
Investigators: Julia P. Baker, Barbara Wanderley, and Danielle O'Brien	
The Winyah Bay Master Naturalist Program: Transforming community members into active stewards of our diverse South Carolina habitats	17
Investigator: Jennifer Plunket	
Blood fluke parasites in the North Inlet estuary: Diversity and life cycles.....	18
Investigators: Isaure de Buron and Dennis Kyle	
Terrestrial response to sea level rise as detected through dendrochronology, geomorphology, and hydrology	18
Investigators: Raymond Torres and Richard Keim	
Experimental varying of the marsh platform and macrophyte response	18
Investigators: James T. Morris and Karen Sundberg	
Within-season patterns of larval demersal fish abundance, age, and growth in tidal creeks	18
Investigators: Juliana M. Harding, Dennis M. Allen, and students	
Novel inexpensive instrumentation for long term monitoring of water levels and ground movement in salt marshes.....	19
Investigators: Vitalii A. Sheremet and Erik M. Smith	
Quantitative descriptions of North Inlet oyster (<i>Crassostrea virginica</i>) population biology	19
Investigator: Juliana M. Harding	
South Carolina Estuarine and Coastal Assessment Program (SCECAP)	19
Investigators: Denise Sanger, Andrew Tweel, Sharleen Johnson, David E. Chestnut, Ed F. Wirth	
Intertidal creek geomorphology on semi-annual and decadal timescales.....	20
Investigators: Alex Gorr, Scott White, Matthew E. Kimball, Erik M. Smith, Dennis M. Allen, and Kyle Houser	
Goby and blenny movements, fidelity, and habitat use	20
Investigators: Juliana M. Harding, Dennis M. Allen, and students	

Tracking gravid female movements of diamondback terrapins (<i>Malaclemys terrapin</i>) in the North Inlet estuary	20
Investigators: Danielle Capella and Scott Parker	
Assessing South Carolina estuaries for contamination affecting fish embryogenesis using the Atlantic silverside (<i>Menidia menidia</i>)	20
Investigator: Shannon Davis	
NERR emergent vegetation bio-monitoring: Effects of sea level on the spatial dynamics of salt marsh vegetation communities in the North Inlet estuary	21
Investigators: Tracy Buck, Erik M. Smith, Julie L. Krask, and Baker Stevens	
Shorebird monitoring in the North Inlet estuary	21
Investigators: Jennifer Plunket and citizen scientist volunteers Wendy Allen and Emma Boyer	
Characterization of oyster cement	21
Investigators: Jonathan Wilker and Paul D. Kenny	
The effects of inhibited carbonic anhydrase on the phytoplankton communities in coastal waters	22
Investigators: Eilea Knotts and James L. Pinckney	
Evaluating intertidal oyster reef restoration success	22
Investigators: Keith Walters, Thomas Funk, and students	
A collaborative science program for the National Estuarine Research Reserve System (NERRS): Working with end users throughout the applied research process	22
Investigators: Dwayne E. Porter, Melissa Ide, and Jeremy Cothran	
Long-term measurements of production and physiological ecology of <i>Spartina alterniflora</i>	23
Investigators: James T. Morris and Karen Sundberg	
Co-evolution of complex traits associated with key innovation: Armor and vision in the snapping shrimp (<i>Alpheus heterochaelis</i>)	23
Investigators: Alexandra C. N. Kingston, Daniel I. Speiser, and Rebecca Lucia	
Long-term monitoring of grass shrimp as a bioindicator of non-point source runoff in South Carolina watersheds	23
Investigators: Peter Key, James Daugomah, and Blaine West	
The National Estuarine Research Reserve System Centralized Data Management Office	24
Investigators: Dwayne E. Porter, Melissa Ide, Jennifer Kesse, Amber Knowles, Brooks Folk, Lee Shutt, Dan Ramage, and William H. Jefferson	
Effect of wrack accumulation on salt marsh vegetation, Baruch Institute, Georgetown County, South Carolina	24
Investigators: Richard Stalter, A. Jung, A. Starosta, John Baden, and M. D. Byer ³	
Estimating above ground biomass utilizing Planet multispectral satellite data	24
Investigators: Gwen Miller and James T. Morris	
Hard clam (<i>Mercenaria mercenaria</i>) population dynamics in North Inlet estuary tidal creeks	25
Investigator: Juliana M. Harding	
Physical characteristics of estuarine waters: Long-term monitoring in the North Inlet and Winyah Bay estuaries	25
Investigators: Erik M. Smith, Baker Stevens, Julie L. Krask, and Tracy Buck	
Mosquitoes of the Carolinas	25
Investigator: Brian Byrd	

Analyzing the seasonal change of heavy metal sediment content in Winyah Bay estuary tidal marshes	25
Investigators: Isabel Fangman and Jennifer Hawk	
Examination of potential juvenile tarpon (<i>Megalops atlanticus</i>) overwintering habitat in the North Inlet estuary	26
Investigators: Matthew E. Kimball, Dennis M. Allen, and volunteers	
Development of models for phytoplankton-nutrient responses in support of numeric nutrient criteria for estuarine water quality	26
Investigators: James L. Pinckney, Erik M. Smith, and Krystyn Kibler	
Benthic foraminifera as indicators of ecosystem health in the North Inlet estuary	26
Investigators: Kelly Gibson and Sarah Shealy	
Satellite observations of marsh dieback events and potential environmental influences along coastal South Carolina, 2000-2017	27
Investigators: Cuizhen (Susan) Wang and Jean T. Ellis	
American alligator (<i>Alligator mississippiensis</i>) movement ecology in coastal South Carolina	27
Thomas R. Rainwater, Abigail J. Lawson, and Patrick G.R. Jodice	
Painted bunting (<i>Passerina ciris</i>) breeding survey	27
Investigators: Jennifer Plunket and citizen scientist volunteers Wendy Allen, Bill Brabson, and Marlene Konsek	
Determining age structure of juvenile tarpon (<i>Megalops atlanticus</i>) in nursery habitats in South Carolina estuaries	28
Investigators: Garrett Elmo, Matthew E. Kimball, and Derek P. Crane	
Relating nonpoint source BOD loading to land-use and stormwater management practices in coastal South Carolina	28
Investigators: Erik M. Smith, Claudia Benitez-Nelson, and Kelly McCabe	
3-D subsurface modeling of a clastic paleo-shoreface deposit	28
Investigator: Patrick Duff	
Satellite observations of marsh dieback events and spatial assessment of environmental influences	29
Investigators: Huixan Li and Cuizhen (Susan) Wang	
Weather and climate measurements: Long-term monitoring at Oyster Landing pier	29
Investigators: Erik M. Smith, Baker Stevens, Julie L. Krask, and Tracy Buck	
Diet of terrestrial-mammalian carnivores that have access to marine-aquatic resources compared to terrestrial upland habitats in coastal South Carolina	29
Investigators: Amanda L. Wilson, David Jachowski, Ernie Wiggers, Michael Childress, Patrick Gerard, and Lisette Waits	
Validation of back-calculation methods for juvenile tarpon (<i>Megalops atlanticus</i>)	30
Investigators: Garrett Elmo ¹ , Derek Crane ¹ , Matthew Kimball ²	
Enhancing bivalve aquaculture through species improvement and diversification	30
Investigators: Ximing Guo, Daphne Munroe, Lisa Calvo, David Bushek, Michael DeLuca, and Pete Rowe	
Chemical analysis of soil samples to detect events in a linear model	30
Investigators: Gabrielle Johnson, Ashley Davis, Samantha McClain, and Students Engaged in Aquatic Sciences members	
Associations between temperature and larval recruitment of a non-native crab (<i>Petrolisthes armatus</i>) near its northern range limit	31
Investigators: Teresa Popp and Dara Wilber	

Tarpon (<i>Megalops atlanticus</i>) recruitment to marsh pools and impoundments along the South Carolina coast	31
Investigators: Garrett Elmo, Derek Crane, Matthew E. Kimball, and Erin Hosto	
<i>Petrolisthes armatus</i> density and performance along a latitudinal gradient	31
Investigator: Jeb Byers	
Archaeological investigations of an early to late 18th century house site - testing methods of intrasite patterning detection.....	32
Investigators: Heathley Johnson and Karen Smith	
Conservation of diamondback terrapin (<i>Malaclemys terrapin</i>) nesting sites in the North Inlet estuary: Using predator vocalization to deter raccoon (<i>Procyon lotor</i>) nest predation.....	32
Investigators: Colleen Naeger and Scott Parker	
Using genetic techniques to understand the mechanism behind the expression of circatidal clocks in an estuarine crab	32
Investigators: Paola López-Duarte, Taylor Parker, and Leyna Pence	
Physiological plasticity and adaptation to microhabitats within populations of the eastern oyster (<i>Crassostrea virginica</i>)	32
Investigators: Denise Furr, Anna Ivanina, and Adam Reitzel	
Characterizing the population genomics of the hemichordate <i>Saccoglossus kowalevskii</i>.....	33
Investigators: Caitlin Redak, Ken Halanych, and Michael Tassia	
Developing a behavioral bioindicator of habitat health for coastal salt marshes	33
Investigator: Blaine D. Griffen	
The effects of climate change on <i>Littoraria irrorata</i> and <i>Ilyanassa obsoleta</i> distribution and movement.....	33
Investigators: Samantha McClain and Students Engaged in Aquatic Sciences members	
Analysis of benthic microalgae composition in salt pan and vegetated marsh sediments in the North Inlet estuary, South Carolina	33
Investigators: Ashley Foulk, Ashley Davis, Abby Davis, and Students Engaged in Aquatic Sciences members	
Geographic distribution of gelatinous zooplankton in the North Inlet-Winyah Bay estuarine system	34
Investigators: Joshua Stone and Imani Haley	
Heart development and molecular characterization of the teleost <i>Megalops atlanticus</i>	34
Investigators: Haley A. Davies, Matthew E. Kimball, Aaron J. Adams, and Frank L. Conlon	
The implications of geographic location and intraspecific size variation on <i>Littoraria irrorata</i> personality types	34
Investigators: Christina Salerno and Stephanie Kamel	
Author Index	35




Figure 1: Map of the North Inlet-Winyah Bay estuarine system in Georgetown County, South Carolina.

Fluorescent dissolved organic matter dynamics in the North Inlet estuary

Investigators: Erik M. Smith, Julie L. Krask, Baker Stevens, Tracy Buck, and Susan Denham
North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

There is growing interest in the use of the inherent optical properties of dissolved organic matter (DOM) as proxies for dissolved organic carbon (DOC) concentrations and biogeochemical cycling in coastal ecosystems. This study employs a fluorescent dissolved organic matter (FDOM) optical probe, recently available as part of Xylem/YSI's EXO water quality sonde, to quantify high-frequency DOC dynamics in the North Inlet estuary. Beginning in August of 2012 an EXO equipped with an FDOM probe, together with temperature, salinity, pH, dissolved oxygen, and turbidity probes, has been deployed at the Oyster Landing long-term monitoring station of the North Inlet–Winyah Bay National Estuarine Research Reserve. Initial results have shown that over the majority of FDOM ranges observed to date, FDOM measures can serve as a reliable proxy for DOC concentration once temperature sensitivities and turbidity interferences are accounted for. Ongoing sampling is being conducted to understand the effects of different dissolved organic matter sources on FDOM – DOC relationships as well as the potential issues associated with sample quenching at high FDOM concentrations. This study will allow the temporal dynamics of DOC, the largest pool of organic carbon in marine waters, to be resolved at frequencies not previously possible.

Ecology, behavior, and population biology of bottlenose dolphins (*Tursiops truncatus*) in the North Inlet estuary and adjacent waters

Investigators: Rob Young and students
Department of Marine Science, Coastal Carolina University (SC)

This long-term project, begun in 1997, has investigated various questions related to the ecology, behavior, and population biology of bottlenose dolphins in the North Inlet and Winyah Bay (NIWB) estuaries. Using photo-ID and focal follow and transect surveys, we have identified long-term resident dolphins in the NIWB system, including several documented residents of over 20 years. NIWB dolphin abundance can exceed 100 in the summer but declines in winter as dolphins spend more time in coastal waters or move into other estuaries. About 12 to 15 dolphins are core users of North Inlet, meaning they are sighted more frequently in the North Inlet salt marsh creeks than in Winyah Bay. These dolphins consume a significant proportion of the prey fish populations (11-14 metric tons per year) and we estimate that 3-7% of the annual primary production in North Inlet estuary is required to support them. Dolphin distribution in North Inlet has been correlated with changing patterns of salinity and prey distribution. Mothers with young calves apparently favor low current areas (tidal nodes), and salt marsh residents swim slower and expend less energy while traveling than coastal dolphins. Bottlenose dolphins are managed under the Marine Mammal Protection Act, and the NIWB dolphins are part of the Northern South Carolina Estuarine System stock. We have observed infrequent movements south to Cape Romain and north along the coast to Murrells Inlet. We are currently investigating environmental DNA (eDNA) as a tool to identify dolphin presence and to potentially correlate eDNA concentration with abundance.

Interspecific competition among some salt marsh perennials in South Carolina

Investigators: Richard Stalter¹ and John Baden²
1 - Department of Biological Sciences, St. John's University (NY)
2 - US Army Corps of Engineers, NC, Retired

Salt marsh vegetation in the United States is characterized by distinct zonation of vascular plants. Zonation is less pronounced in brackish versus high salinity marshes. Previous transplant experiments indicated several species could not tolerate conditions in areas where they are not normally found. These experiments, however, failed to differentiate the effects of abiotic and biotic (namely interspecific competition) factors. Controlled, reciprocal transplant manipulations have been performed. Growth and survival were monitored to measure the relative importance of interspecific competition and abiotic factors as determinants of zonation patterns between the salt marsh cord grass, *Spartina alterniflora*, and the black needle rush, *Juncus roemerianus*. *Spartina alterniflora* was able to invade the *J. roemerianus* zone when the latter was removed from land that it originally occupied in the marsh. *Juncus* marginally invaded the *S. alterniflora* zone when the latter was removed. *Juncus* did not transplant well; almost 100 % of the transplanted *J. roemerianus* died even when dug up and replanted in place.

Construction of biodegradable floating treatment wetland utilizing wood and bamboo

Investigators: William Strosnider¹, Travis Tasker², Lily Currie², and Kendra Zaruba²

1 - Baruch Marine Field Laboratory, University of South Carolina

2 - Department of Environmental Engineering, Saint Francis University

The purpose of this research was to develop new designs for floating treatment wetlands (FTWs) that utilize natural and biodegradable materials to support plant growth and improve water quality. In this work, we were able to design biodegradable and durable wood and bamboo floating rafts embedded with coir fiber mats. While the rafts were still floating after 4 weeks and were planted with *Pontederia cordata*, the rafts need to be continually monitored in the future to determine the longevity of the floating mats. The benefits and success of constructing FTWs out of more natural, biodegradable, and ubiquitous materials could reduce the costs of FTWs and allow for more resilient plant and biological communities.

A forty-seven year comparison of the vascular flora at three abandoned rice fields, Georgetown, South Carolina

Investigators: Richard Stalter¹, Joseph Rachlin², and John Baden³

1 - Department of Biological Sciences, St. John's University (NY)

2 - Lehman College (NY)

3 - US Army Corps of Engineers, Retired

The vascular flora present in three abandoned rice fields of the Winyah Bay estuary at the Belle W. Baruch Institute for Marine and Coastal Sciences, Georgetown County, South Carolina identified in 1968-1969 was compared with the vascular flora present in 1987-1991, and 2013-2015. Twenty vascular plant species were identified in 1968-1969 and 22 in 2013-2015 at the most saline marsh, Thousand Acre Rice Field. Forty seven taxa were reported at Airport marsh in 1968-1969 and 27 in 2013-2015. Fifty six taxa were reported at Alderly marsh in 1968-1969, while only 41 were identified here in 2013-2015. A parsimony algorithm was used to evaluate the distribution and co-occurrence of vascular brackish marsh species in 3 abandoned rice fields sampled at three intervals, 1968-1969, 1987-1991, and 2013- 2015. There was a shift in the flora at the two least saline sites, Alderly marsh and Airport marsh from 1968-1969 to 1987-1991 and 2013-2015. Three factors, rising sea level, an increase in water salinity plus the invasion of *Phragmites australis* may explain the shift in vegetation in Alderly marsh and Airport marsh, the two least saline marshes. There was a shift in the flora at the most saline site, Thousand Acre Rice Field from 1967-1969 to 1987-1991 and 2013-2015 after the marsh was savaged by Hurricane Hugo in 1989. The invasion of non-native *Phragmites australis* at all sites and the increase in water salinity at all sites best explains the reduction in vascular plant species at Airport and Alderly marshes over the 47-year collecting period.

Fish and crustacean use of marshes and intertidal creeks: Population and community level changes and relationships with weather and climate-driven changes

Investigators: Bruce W. Pfirrmann, Matthew E. Kimball, Dennis M. Allen, and Paul D. Kenny

Baruch Marine Field Laboratory, University of South Carolina

Collections of nekton (fishes, shrimps, and crabs) have been made in the Oyster Landing marsh-creek basin from 1984 to the present. The objective has been to track the composition, abundance, biomass, and length distributions of nekton and determine patterns, trends, and factors influencing changes over seasons, years, and decades. From 1984-2003, this effort was based on biweekly seine hauls from an isolated pool (low tide) in the intertidal creek. In 1996, we started a new time series from the flooded marsh surface (high tide) adjacent to the creek. From 1996-2003, both the low tide seine and high tide enclosure collections were made on the same day and tide. These efforts demonstrated that although patterns of occupancy of the intertidal basin by species and life stages within species varied, they were predictable according to tide level and time of year. Large, short-term changes in salinity and temperature affected abundance in the intertidal basin, but familiar patterns were quickly re-established. High tide enclosure collections continued through 2011. In 2012 the sample site shifted to an adjacent isolated section of creek at low tide, where from 2012-2018, sampling focused on documenting shifts in the timing, size, and growth of juvenile transient species. Beginning in 2019, our effort expanded to include the entire nekton assemblage occurring in the tidal creek at low tide. The long-term time series is increasingly important as we interpret impacts of global climate change on nekton populations and the shallow water habitats that are essential to their development. The results are used to inform the management of salt marsh-estuaries, watersheds, and fisheries in the southeastern US.

The conservation status of the canebrake rattlesnake (*Crotalus horridus atricaudatus*) at Hobcaw Barony, with identification of key areas for conservation of its herpetofauna

Investigator: Allan L. Markezich

Department of Natural Sciences, Black Hawk College (IL)

This ongoing long-term project assesses the abundance and microgeographic and ecological correlations of canebrake rattlesnake (*Crotalus horridus atricaudatus*) occurrence in the approximate 8,000 acres of terrestrial communities of Hobcaw Barony. Observations involve timed road and walking surveys and usage of drift fences and cover boards along with various marking techniques of specimens to assess abundance. Data taken on snake occurrences involve coordinates of specific geographic localities, topography, general and specific ecological characteristics of communities, and variables involving specific microhabitat and seasonal associations. Data taken by others and information on historical land usage at Hobcaw Barony are also utilized. Results currently indicate that a relatively small metapopulation of the canebrake rattlesnake exists on the property, with highest densities in specific and relatively small areas. Hardwood forests and palmetto swamplands bordering upland areas are key ecological components of this species' environment at Hobcaw Barony. The study to date indicates that the current conservation status of the canebrake rattlesnake on the property is poor, and that populations may have declined in the past five years. Management efforts should be made to conserve critical habitats on the property and to minimize disturbance of them. People using vehicles should also be vigilant of snakes on the roads to reduce road mortality of individuals, which has increased in the past several years. Similar information on occurrence and abundance of other reptilian and amphibian species is also used to understand geographic and habitat correlates of herpetofaunal diversity (i.e., species richness) on the property. Hardwood forests and freshwater wetlands, and associated ecotones between these and pine forests, are critical areas for herpetofaunal diversity, with the greatest species richness found in the northern portion of the property. Managed pine forests have had the lowest richness.

Seasonal energetics of the blue crab (*Callinectes sapidus*)

Investigators: Blaine D. Griffen¹, Matthew E. Kimball², and Bruce W. Pfirmann²

1 - Department of Biology, Brigham Young University

2 - Baruch Marine Field Laboratory, University of South Carolina

The goal of this project is to examine seasonal changes in blue crab individual energetics across age classes and genders. The primary methods used involve physiological analysis of the gonads and hepatopancreas to determine seasonal patterns in size and lipid composition of these organs. This will be used to determine patterns in long-term energy storage and energy allocation towards reproduction. In addition, short-term energy reserves will be assessed using glycogen content of muscle tissues. This information will then be combined with bioenergetics growth models for this species to understand seasonal patterns in the energetics of growth and reproduction.

Chemical characteristics of estuarine waters: Long-term monitoring at four sites in |North Inlet estuary

Investigators: Erik M. Smith, Julie L. Krask, Baker Stevens, and Susan Denham

North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

As part of the National Estuarine Research Reserve System (NERRS) System-Wide Monitoring Program (SWMP), water chemistry sampling was initiated in June 1993 to monitor concentrations of suspended solids, total nitrogen, ammonium, nitrate, nitrite, total phosphorus, orthophosphate, and chlorophyll *a* at four locations within the North Inlet–Winyah Bay NERR. Water samples are collected every 20 days with ISCO automated water sampling devices at intervals of 2 hours and 4 minutes over two complete tidal cycles. Sampling and chemical analyses adhere to strict national protocols developed as part of the NERRS System-Wide Monitoring Program. The consistent, long-term collection of water chemistry variables allows for the characterization of short-term variability and detection of long-term change in key water quality parameters. These data also provide critical information for various studies of biological and physical processes in the North Inlet estuary. Data, along with detailed metadata, are sent to the NERRS Centralized Data Management Office (CDMO) for quality assurance and quality control, and then made available via the CDMO website (<http://cdmo.baruch.sc.edu>). Water chemistry data collected in North Inlet prior to the initiation of the NERRS SWMP sampling (some dating back to 1978) are available via the Baruch website's Data and Publications link (<http://www.baruch.sc.edu/water-quality-chemistry-databases>).

Maintenance and operation of IOOS/SECOORA priority WERA HF radar sites

Investigators: George Voulgaris¹ and William H. Jefferson²

1 - School of the Earth, Ocean, and Environment, University of South Carolina

2 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

The objective of this study is to remotely monitor the ocean surface currents and waves in Long Bay using two high frequency (HF) radar stations. Scientists from the University of South Carolina operate and maintain two US IOOS/SECOORA identified priority WERA system radar sites (Georgetown, SC and Fort Caswell, NC). One station is located on Hobcaw Barony (33°21'19.60"N, 79° 9'12.56"W) and the other station is located at Caswell Beach, NC (33°53'25.18"N, 78° 1'40.64"W). Each station remotely measures the surface ocean currents up to 120 miles offshore and when combined, create maps of temporal and spatial distribution of waves and currents over the entire Long Bay area. Data from these sites are sent to SECOORA and National High Frequency Radar Network for integration, display, and dissemination.

Painted Bunting (*Passerina ciris*) monitoring project

Investigators: Jennifer Plunket¹ and citizen science volunteers Wendy Allen, Bill Brabson, Cindy Kerr, Marlene Konsek, Pete Little, and Sandy Little

1 - North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

Painted Buntings (PABUs) are the most colorful of the migratory songbirds that visit the coast of South Carolina. Adult males sport a royal blue head, neon green back, and red breast and rump. PABUs return to the area mid-April, nest in shrubs near marshes, and migrate south in the fall to central and southern Florida, Cuba, and the Yucatan peninsula of Mexico. Surveys conducted since 1966 have demonstrated a decline in PABUs which lead to the establishment of a monitoring project that includes banding at select sites in the southeast and observations made by citizen scientists. This project, Painted Bunting Observation Team or PBOT, was headed up by scientists at UNC Wilmington. The North Inlet–Winyah Bay National Estuarine Research Reserve (NI-WB NERR) served as a PBOT banding site 2007-2012, and has maintained a feeder near the Baruch Marine Field Laboratory (BMFL) since 2007. The NI-WB NERR established a Painted Bunting monitoring project in summer 2014 to document buntings coming to the feeder. Reserve staff and citizen scientists make timed observations of PABUs and record color band combinations, if present. Citizen scientists continue to work with Reserve staff to monitor PABUs on Hobcaw Barony during the 2019 nesting season. The project will contribute to a better understanding of PABU longevity, how long they stay during the nesting season, whether they come back to the same site from year to year, and other aspects of their natural history.

Eddy covariance flux measurements to quantify salt marsh productivity and its response to environmental variability over multiple time scales

Investigators: Thomas L. O'Halloran¹ and Erik M. Smith²

1 - Baruch Institute for Coastal Ecology and Forest Science, Clemson University (SC)

2 - North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

Accurate and integrative measures of marsh productivity as well as the sensitivity of marsh production to environmental variability over multiple time-scales are essential to understanding how salt marshes will respond to future environmental and anthropogenic stressors. This study is employing state-of-the-art eddy covariance flux instrumentation (IRGASON, Campbell Scientific) to generate high-frequency (30-minute interval) measurements of terrestrial-atmospheric CO₂ exchange at spatial scales large enough (on the order of 20,000 m²) to capture landscape-level dynamics. The instrumentation is located with the National Estuarine Research Reserve's existing salt marsh monitoring infrastructure within the Crabhaul Creek marsh of the North Inlet estuary to leverage ongoing data collection of marsh vegetation, surface elevation and tidal inundation, salinity and pore water chemistry, and meteorological data. Results of this study will greatly improve our understanding of marsh sensitivity to environmental variability and change through the development of empirical models relating the integrated response of the salt marsh ecosystem (as gross primary production, ecosystem respiration, and net ecosystem exchange) to environmental variability over temporal scales not previously possible and at spatial scales large enough to integrate landscape-level responses.

Benthic bivalves as potential indicator species for ecosystem climate change effects

Investigators: Juliana M. Harding¹ and Dennis M. Allen²

1 - Department of Marine Science, Coastal Carolina University (SC)

2 - Baruch Marine Field Laboratory, University of South Carolina

The current project builds on the Baruch Institute's long-term monitoring program describing macrobenthic bivalve trends in the North Inlet estuary. The coupling of modern and historic data allows evaluation of potential changes in species richness and recruitment timing since 1982 related to increasing winter water temperatures. Modern field collections (Bly Creek, Bread and Butter Creek) describing the current status of macroinfauna will be coupled with historic macroinfaunal sample analyses to 1) quantitatively describe infaunal bivalve populations and demographics, and 2) evaluate the potential for increasing water temperatures to change these dynamics and related ecosystem services over decadal scales. This research complements previous and ongoing work describing other North Inlet ecosystem trophic levels and will enhance existing descriptions and predictions of ecosystem function.

Coastal training activities in the North Inlet–Winyah Bay National Estuarine Research Reserve: Protecting water and habitat quality through science-based community training

Investigator: Maeve Snyder

North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

Coastal training activities connect local decision makers to the emerging research and scientific knowledge generated to help the decision makers make more informed decisions on coastal environmental issues. The Coastal Training Program provides needs-based workshops, trainings, and tools to decision makers in Georgetown and Horry counties and these efforts especially target county and municipal staff and officials, and those decision makers that strongly influence local land use, such as planners, developers, engineers, and realtors, as well as those with a role in natural resource management within local counties and municipalities. The Coastal Training Program works to protect water and habitat quality in a region of rapidly developing coastal communities by providing science-based training events on the issues of stormwater management and low impact development principles, habitat protection and restoration, coastal hazards and climate change, and other emerging priority issues. The program frequently partners with the ACE Basin National Estuarine Research Reserve, South Carolina Sea Grant Consortium, the Coastal Waccamaw Stormwater Education Consortium, the Clemson University Extension Service, and the Carolina Clear Program.

Toxicological effects of tire wear particles on fathead minnow (*Pimephales promelas*) and Atlantic killifish (*Fundulus heteroclitus*)

Investigators: Peter van den Hurk¹, Stephanie LaPlaca², and John Weinstein³

1 - Department of Biological Sciences, Clemson University (SC)

2 - Environmental Toxicology Program, Clemson University (SC)

3 - Department of Biology, The Citadel (SC)

Recent studies on the distribution of microplastics in the Charleston Harbor, South Carolina, revealed that a large part of the microplastic particles that are found in the intertidal sediments are tire wear particles. These particles originate from the wear of tire treads on roadways, and wash into the estuary during rain events. The abundance of these particles has raised questions about potential toxicity to aquatic organisms that may ingest these particles. To investigate the potential toxicity of tire wear particles we started a project comparing the effects in both fathead minnow and Atlantic killifish. The fish are exposed to different concentrations of tire wear particles in a 7 day exposure. Earlier experiments in fathead minnow revealed that particles were actually ingested and accumulated in the intestinal tract. At the highest concentration tested (6000 mg/l) we also observed partial mortality in the fathead minnow, which is therefore close to the LC50. To investigate if polynuclear aromatic hydrocarbons were leaching from the particles, bile fluorescence is measured, together with potential induction of cytochrome P450-1A through the EROD assay. In addition, glutathione S-transferase is measured as a general stress parameter. The results of this study will allow a comparison between the two species, with the consideration that Atlantic killifish is an estuarine species that is much more used to being exposed to high particle load in its natural environment than the fathead minnow. The results of the project may be used to support environmental management strategies that reduce particulate matter in road runoff from entering estuarine environments.

Public and K-12 education activities: North Inlet–Winyah Bay National Estuarine Research Reserve

Investigators: Beth Thomas and Hayley Fournier

North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

Educational activities for the general public, K-12 teachers, and students highlighting coastal ecology and integrating findings from research are offered throughout the year. Seasonal schedules of public outreach activities are produced throughout the year, and programs are promoted through informational fliers, newsletters, newspapers, and website (www.northinlet.sc.edu), and the Reserve's Facebook page in addition to local online community event calendars. Program offerings include estuarine and beach ecology activities for all ages, biking and kayaking programs featuring coastal ecology, open houses and research lectures, and research-based citizen science programs. Professional Teacher Development opportunities and field trips for K-12 public, private, and homeschool students are also available, as well as job shadowing and research experiences for middle and high school students. Off-site outreach includes presentations to environmental and civic groups, local festivals, special outreach programs at regional libraries and museums, afterschool programs for local elementary and middle schools, science and environmental fairs, and career days. Partnerships with other local environmental education providers, including the Belle W. Baruch Foundation, ACE Basin National Estuarine Research Reserve, South Carolina Department of Natural Resources, South Carolina Sea Grant Consortium, Friends of Coastal South Carolina, the Waccamaw National Wildlife Refuge, and the Coastal Waccamaw Stormwater Education Consortium provide additional opportunities for public education, teacher training, and professional development, as well as staff and resources for enhanced programming and outreach.

Saltwater intrusion monitoring

Investigators: Alicia Wilson¹ and William Clendenin²

1 - School of the Earth, Ocean, and Environment, University of South Carolina

2 - South Carolina Department of Natural Resources

Knowledge of the location of the freshwater-saltwater interface in coastal aquifers is critical for managing coastal groundwater resources, for predicting saltwater intrusion, and for calculating groundwater-related chemical exchange between aquifers and the coastal ocean. This project installed permanent wells to monitor salinity and saltwater intrusion in the upper (up to 100 ft depth) aquifers at the North Inlet estuary, as part of the South Carolina Department of Natural Resources long-term coastal monitoring network. To date, the stratigraphy from the well logs has been used to support the development of regional groundwater flow models. The models are further constrained by hydraulic head and salinity observations from the wells. Monitoring is ongoing.

Long-term changes in the zooplankton of the North Inlet estuary and relationships with climate change

Investigators: Dennis M. Allen¹, Josh Stone², Bruce W. Pfirrmann¹, and Matthew E. Kimball¹

1 - Baruch Marine Field Laboratory, University of South Carolina

2 - Department of Biological Sciences, University of South Carolina

Collections have been made at the same location, stage of tide, and time of day every two weeks since 1981. Oblique tows with 153 μm mesh nets collect copepod and small invertebrate larvae, and 365 μm epibenthic sled tows capture larval fishes, shrimps, and crabs and other large zooplankton species. Seasonal and interannual changes in abundance, diversity, and species composition of the assemblages in Town Creek are documented and related to fluctuations and trends in the physical characteristics of the estuary. Information is collected for more than 50 taxonomic groups and species which are indicators of the condition and diversity of life in the estuary. Constituents include species of ecological and economic importance to the southeastern US region. Large, long-term decreases in the abundance of small (153 micron) zooplankton and changes in the timing of occurrences of some larval fishes and decapods (365 micron) have indicated responses to a long-term increase in water temperature. We have also observed responses of zooplankton populations to climatic events including El Niño-Southern Oscillation, tropical storm-related reductions in salinity, and drought. The value of these datasets continues to increase as we formulate and test new hypotheses about impacts of climate change on estuarine ecosystems and fisheries.

Quantifying elevational and latitudinal gradients in marsh snail predation

Investigators: Rebecca Atkins, Craig Osenberg, and Daniel Hawkins
Odum School of Ecology, University of Georgia

In southeastern salt marshes, the periwinkle snail (*Littoraria littoraria*) is a dominant grazer, feeding on stalks of smooth cordgrass (*Spartina alterniflora*). Previous work has demonstrated that *Littoraria*'s effect on *Spartina* depends on the abundance and size structure of its population. *Littoraria* body size and abundance is known to vary across a latitudinal gradient from Florida to Maryland, but we do not know the cause of this variation. *Littoraria* are eaten by mud crabs, blue crabs, birds, and terrapin turtles. However, little is known about the role that these predators play in determining *Littoraria* population size structure and abundance. To gain additional information, "tethering experiments" will be conducted during the summer in the North Inlet estuary as well as other marsh sites from Florida to Maryland.

Mechanisms for thermal tolerance in an estuarine cnidarian

Investigator: Adam Reitzel
Department of Biological Sciences, University of North Carolina Charlotte

Estuarine species like the starlet sea anemone, *Nematostella vectensis*, experience large fluctuations in temperature, requiring wide temperature tolerances, and at the same time, have a natural distribution along a pronounced thermal cline (Atlantic coast of North America), which may promote the evolution of different temperature optima and tolerances in populations. Previous research with *Nematostella* has shown extensive genetic variation, some of which is strongly segregated between populations, and large differences in phenotypic variation in response to both acute and chronic temperature, both of which vary with site of collection, consistent with adaptation to temperature. The North Inlet–Winyah Bay estuarine system represents the most southern location where successful collections for this species have occurred in recent decades and individuals from this site appear to have elevated thermal tolerance. We are using field-deployed temperature loggers coupled with seasonal field samples to compare mean and maximum temperatures with the growth and gene expression for this species. Results will be compared with *N. vectensis* collected from sites as far north as Nova Scotia to understand the genetic processes that allow these estuarine specialists to adapt and respond to their dynamic temperature environments.

Green porcelain crab (*Petrolisthes armatus*) larval biology and phenology

Investigator: Juliana M. Harding
Department of Marine Science, Coastal Carolina University (SC)

Green porcelain crab (*Petrolisthes armatus*) larval biology and phenology will be described with field collections. Weekly zooplankton tows will be used to monitor the presence and stages of crab larvae. Larval morphology will be described from field-caught specimens. This information will add to the understanding of this invasive (extended geographic range) species' occurrence in the North Inlet estuary. *Petrolisthes armatus* larvae first occurred in biweekly zooplankton collections in the mid-1990s and little is known about the timing, periodicity, and duration of larval production. Modern patterns in larval occurrence can be compared to similar information from archived biweekly collections to evaluate potential changes with temperature.

Exploring macroinvertebrate diversity of non-native macroalgae *Agarophyton vermiculophyllum*

Investigator: Timothy Lee
Department of Biology, East Carolina University (NC)

The major goal of this study is to understand the macroinvertebrate diversity associated with the non-native macroalgae red seaweed (*Agarophyton vermiculophyllum*), which is native to the northwestern Pacific. My study in South Carolina is a small subset of the larger biogeographic study on macroinvertebrate diversity, so my sampling site in the North Inlet estuary is just one of the many sites that I plan on sampling along the US east coast. For field work, I will collect five replicates of seaweed. Macroinvertebrates from each replicate of seaweed will be extracted for species identification. My study at the North Inlet estuary will help answer the bigger question of macroinvertebrate diversity in estuarine habitats transformed by non-native seaweed.

Sediment accretion in North Inlet estuary salt marshes

Investigators: James T. Morris¹ and Karen Sundberg²

1 - Department of Biological Sciences, University of South Carolina

2 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

The objective of this study is to understand how the elevation of the marsh surface is regulated. A major hypothesis being tested is that eutrophication initiates a sequence of changes in the sediments, beginning with a decrease in volume due to enhanced decomposition of organic matter. In fact, sediment accretion in experimentally fertilized marsh plots has increased. This is probably due to an increase in sedimentation caused by a higher density of plant stems in fertilized plots. Marsh plots were fertilized from 1996 or 2001 until 2004. A surface elevation table (SET) is used to measure marsh elevations in low and high marsh *Spartina alterniflora* plots approximately monthly. Currently we are looking at the effect of decreasing eutrophication on marsh surface elevation, and we hypothesize that there will be a decrease in volume of below ground biomass due to enhanced decomposition now that below ground production is no longer stimulated. Results of a model linking plant production and sedimentation with sea level indicate that the marsh maintains its elevation with respect to mean sea level for a range of rates of sea-level rise, up to a threshold. The elevation of the marsh platform with respect to mean sea level is inversely proportional to the rate of sea level rise.

Analyzing nickel, zinc, and lead in *Fundulus heteroclitus*

Investigators: Julia P. Baker, Barbara Wanderley, and Danielle O'Brien

Division of Business, Mathematics, and Sciences, Columbia College (SC)

The zinc, nickel, and lead concentrations in *Fundulus heteroclitus* were assessed on fish from the North Inlet-Winyah Bay estuarine system (NI) and Georgetown Harbor (GH). *Fundulus heteroclitus* from both environments were obtained using minnow traps. Samples from Georgetown Harbor were difficult to obtain, so in addition to the few fish caught, samples previously obtained researchers at Columbia College were also analyzed. However, the livers had been removed from these fish raising another research question on the importance of the liver in metal accumulation in *F. heteroclitus*. Samples, therefore, were analyzed in two categories: with and without livers. The samples were prepared in triplicate, ashed, digested with nitric acid and analyzed twice by atomic absorption spectroscopy. The NI sample without livers had the following metal concentrations measured in ug/g of dry weight: zinc 890 ± 58 $\mu\text{g/g}$, nickel 3.3 ± 0.4 $\mu\text{g/g}$, lead 4.5 ± 0.6 $\mu\text{g/g}$. For the GH sample without livers the metal concentrations were zinc 1090 ± 10 $\mu\text{g/g}$ and nickel 4.4 ± 0.5 $\mu\text{g/g}$. The metal concentrations of the NI sample with livers were zinc 845 ± 20 $\mu\text{g/g}$, nickel 9 ± 3 $\mu\text{g/g}$, and lead 3.9 ± 0.7 $\mu\text{g/g}$. The metal concentrations of the GH sample with livers were zinc 462 ± 69 $\mu\text{g/g}$, and nickel 9.5 ± 5 $\mu\text{g/g}$. The GH samples were not analyzed for lead. The results indicate the liver is important in the accumulation of nickel in *F. heteroclitus* but this correlation was absent for zinc and lead. No clear difference was indicated in the levels of zinc between the NI and GH samples, but the GH samples showed twice as much nickel as the NI samples.

The Winyah Bay Master Naturalist Program: Transforming community members into active stewards of our diverse South Carolina habitats

Investigator: Jennifer Plunket

North Inlet-Winyah Bay National Estuarine Research Reserve, University of South Carolina

The Winyah Master Naturalist course is designed to train community members to become active volunteer stewards of our coastal environment. Participants gain skills in nature interpretation, research methods, and resource protection through 12 day-long field classes occurring on Fridays from March to June. The course involves field trips with expert interpreters to the mountains, forests, swamps and marshes that make South Carolina a unique and beautiful classroom for the nature enthusiast. Students will learn to 'read' the landscape through developing an understanding of the geology, ecology and human impacts on natural habitats. Participants completing the course and 30 hours of approved volunteer work will receive a Master Naturalist certification and will be eligible to join a local chapter and participate in advanced volunteer training courses. Participants do not need to have a background in the natural sciences; a diversity of backgrounds, skills and interests is welcomed.

Blood fluke parasites in the North Inlet estuary: Diversity and life cycles

Investigators: Isaure de Buron¹ and Dennis Kyle²

1 - Department of Biology, College of Charleston (SC)

2 - Department of Global Health, University of South Florida

Blood flukes are pathogenic parasites that infect the cardiovascular system of their fish hosts. Their life cycle uses annelids as intermediate hosts. The objectives of this project are 1) to determine which species of fluke infects spotted seatrout (*Cynoscion nebulosus*) in the North Inlet estuary compared to other South Carolina estuaries and 2) to sample annelids regularly throughout the year in order to unravel the specifics of the life cycles of these particular parasites. This study will also allow us to determine a potential seasonal pattern of infection by the flukes in seatrout and will give us insight into the diversity of blood flukes in the North Inlet estuary.

Terrestrial response to sea level rise as detected through dendrochronology, geomorphology, and hydrology

Investigators: Raymond Torres¹ and Richard Keim²

1 - School of the Earth, Ocean, and Environment, University of South Carolina

2 - School of Renewable Natural Resources, Louisiana State University

We propose to evaluate the rates of salt marsh advance into the terrestrial landscape using dendrochronology. The corresponding tree ring chronology will be used to assess rates of geomorphic change of the terrestrial and intertidal landscapes and the patterns and dynamics of surface and near surface freshwater and salt water flows.

Experimental varying of the marsh platform and macrophyte response

Investigators: James T. Morris¹ and Karen Sundberg²

1 - Department of Biological Sciences, University of South Carolina

2 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

The objective of this study was to design a simple experiment to investigate how varying the marsh platform in relation to mean sea level would affect macrophyte production, stand dynamics, and biomass allocation patterns of salt marsh plants. One specific goal was to ascertain above ground and below ground allocation patterns and quantify where the bulk of below ground biomass was located in relation to marsh elevation and sea level. The experiments were initiated in 2003. Currently there are three marsh planters ('marsh organs'), each with six treatment platform levels that span the upper half of the tidal range, and six replicates per treatment. In general, the marsh organs are planted in March with salt marsh plugs (currently *Spartina alterniflora*) collected nearby; stem height measurements are obtained monthly as an estimate of standing biomass; and plants are harvested at the end of the growing season, to determine above ground and below ground productivity. In recent years, replicates have been selectively harvested such that we now have an age treatment in addition to the elevation treatment. The frequency of inundation results in significant variation in stand densities and plant heights, and we are observing different biomass allocation patterns with time. These changes in stand densities and macrophyte morphology may have profound effects on the ability of salt marshes to accrete allochthonous sediments and maintain pace with sea-level rise. Furthermore, allocation patterns may ultimately influence net annual primary productivity within salt marshes.

Within-season patterns of larval demersal fish abundance, age, and growth in tidal creeks

Investigators: Juliana M. Harding¹, Dennis M. Allen², and students

1 - Department of Marine Science, Coastal Carolina University (SC)

2 - Baruch Marine Field Laboratory, University of South Carolina

Abundance, age, and growth patterns of demersal oyster reef fish larvae including the naked goby (*Gobiosoma bosc*), crested blenny (*Hypoleurochilus geminatus*), feather blenny (*Hypsoblennius hentz*), freckled blenny (*Hyposblennius ionthas*), and striped blenny (*Chasmodes bosquianus*) are being examined. Regular ichthyoplankton collections will be used to describe larval fish abundance and demographics. Fish otoliths will be used to describe age and growth rates. These data will be used in combination with information about goby and blenny larvae cultured at known conditions during 2012 and 2013 to interpret patterns observed in the long-term zooplankton series (1981-present).

Novel inexpensive instrumentation for long term monitoring of water levels and ground movement in salt marshes

Investigators: Vitalii A. Sheremet¹ and Erik M. Smith²

1 - Northeast Fisheries Science Center, National Oceanic and Atmospheric Administration (MA)

2 - North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

An array of six inexpensive tidal gauges and six ground movement detectors is proposed to be deployed at the North Inlet-Winyah Bay National Estuarine Research Reserve. We would like to expand our recent observations at the Waquoit Bay National Estuarine Research Reserve and contrast the effects of different marsh morphology and much larger tidal range. The question is whether the sediment accretion in the marsh is rapid enough to keep up with the sea level rise caused by the climate change. Several Surface Elevation Tables (SET) were established at the Waquoit Bay Reserve and other Reserves to monitor the long term evolution. Recently, we realized that the ground movement caused by tide flooding the marsh needs to be taken into the account. Both the tide gauges and the ground movement detectors are designed by the PI for the project and are based on inexpensive stock Onset Computer Accelerometer Loggers. We propose to make a pilot deployment for a period of 1-2 months in order to record one complete tidal cycle with 1 minute sampling intervals. The observations will allow us to document the propagation of tides within the reserve and corresponding expansion of the top surface of the marsh. The ultimate goal is to develop an observation protocol that could be applied throughout the whole National Estuarine Research Reserve System.

Quantitative descriptions of North Inlet oyster (*Crassostrea virginica*) population biology

Investigator: Juliana M. Harding

Department of Marine Science, Coastal Carolina University (SC)

Oyster (*Crassostrea virginica*) population biology sets the foundation for maintenance and persistence of the biogenic habitat as well as the associated trophic communities and ecological services. These dynamics respond to a variety of factors functioning at time scales ranging from days to decades. This research describes basic oyster population parameters including recruitment intensity and periodicity as well as density, demographics, disarticulation rates, Dermo prevalence and intensity, biomass, and reef spatial extent at sentinel sites in the Town, Clambank, Crabhaul, Debidue, and Bly Creek basins. Environmental data will be collected concurrently and integrated with the biological data. The integrated data sets will be examined in the context of available historic data and documented environmental changes across decadal time scales.

South Carolina Estuarine and Coastal Assessment Program (SCECAP)

Investigators: Denise Sanger¹, Andrew Tweel¹, Sharleen Johnson¹, David E. Chestnut², Ed F. Wirth³

1 - South Carolina Department of Natural Resources

2 - South Carolina Department of Health and Environmental Control

3 - Southeast Regional Office, National Oceanic and Atmospheric Administration (SC)

The South Carolina Department of Natural Resources (SCDNR) and the South Carolina Department of Health and Environmental Control (SCDHEC) have been conducting an ongoing comprehensive collaborative coastal monitoring program (South Carolina Estuarine and Coastal Assessment Program; SCECAP) since 1999. The goal of SCECAP is to annually monitor the condition of the state's estuarine habitats and associated biological resources. SCECAP integrates measures of water and sediment quality with multiple measures of biological condition at a large number of sites throughout the state's coastal zone. It also expands historical monitoring activities that have primarily focused on open water habitats (e.g., bays, sounds, tidal rivers) to include an assessment of conditions in tidal creeks, which serve as important nursery habitat for many species. The SCECAP program, combined with the other cooperating programs, provides a number of benefits including 1) the ability to identify areas of estuarine habitat that are impaired or degraded with respect to a suite of sensitive biological, chemical, and physical measures; 2) a cost-effective standardized protocol that is used by both SCDNR and SCDHEC that is consistent with protocols used in other US coastal states, thus allowing better regional prioritization of stressors and impacts; 3) more comprehensive periodic reports on the condition of water quality and habitat condition throughout the state's coastal zone than could be accomplished by the individual programs alone. As of August 2019, over 840 sites have been sampled statewide, with 11 located in the North Inlet estuary and an additional 37 stations located in the adjacent Winyah Bay.

Intertidal creek geomorphology on semi-annual and decadal timescales

Investigators: Alex Gorr¹, Scott White¹, Matthew E. Kimball², Erik M. Smith³, Dennis M. Allen², and Kyle Houser²

1 - School of the Earth, Ocean, and Environment, University of South Carolina

2 - Baruch Marine Field Laboratory, University of South Carolina

3 - North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

Previous work in 1997-1999 examining nekton use in 8 intertidal creek tributaries of Clambank Creek revealed variations in geomorphology (e.g., depth, bank steepness, etc.) correlated to abundance of resident and transient nekton. The nature of these patterns suggests nekton actively select creeks with preferred geomorphic characteristics, with implications for essential fish habitat, conservation, and restoration efforts. To examine the stability of the intertidal creek geomorphology and the consistency of nekton habitat preferences, the geomorphology of these same 8 intertidal creeks was measured again 15 or more years later. Geomorphological characteristics originally measured in 1997 using traditional surveying techniques were re-measured in 2016 and 2017 using terrestrial laser scanning and 2018 to present using a UAV quadcopter. Digital terrain models derived from these data allow comparison of small seasonal changes within creeks and large, decadal changes among creeks. The goal of this study is to determine if and how much the geomorphology of these creeks has changed over the years.

Goby and blenny movements, fidelity, and habitat use

Investigators: Juliana M. Harding¹, Dennis M. Allen², and students

1 - Department of Marine Science, Coastal Carolina University (SC)

2 - Baruch Marine Field Laboratory, University of South Carolina

Habitat use patterns of demersal oyster reef fishes including naked goby (*Gobiosoma bosc*) crested blenny (*Hypleurochilus geminatus*), feather blenny (*Hypsoblennius hentz*), freckled blenny (*Hyposblennius ionthas*), and Striped Blenny (*Chasmodes bosquianus*) in Crabhaul Creek in the North Inlet estuary are being examined. Artificial nesting substrates have been and will continue to be used to describe movement and fidelity patterns of these resident fishes. Regular surveys will provide information on site fidelity and home range as well as demographics, nest substrate preferences, and habitat use patterns of resident fishes.

Tracking gravid female movements of diamondback terrapins (*Malaclemys terrapin*) in the North Inlet estuary

Investigators: Danielle Capella and Scott Parker

Department of Biology, Coastal Carolina University (SC)

The purpose of this research is to quantify large-scale and small-scale movement patterns and habitat use of diamondback terrapins during the summer activity season. Specifically, we aim to determine if movement and habitat of gravid females differs from males and non-gravid females. This information will help inform future diamondback terrapin conservation efforts.

Assessing South Carolina estuaries for contamination affecting fish embryogenesis using the Atlantic silverside (*Menidia menidia*)

Investigator: Shannon Davis

Department of Biological Sciences, University of South Carolina

Estuarine environments are often called the “nurseries of the sea” because of their importance in providing a protected environment for fish, shellfish, and marine mammal reproduction. They also filter contaminants from fresh water sources, which could potentially impact fish embryogenesis. The objective of my proposal is to characterize the Atlantic silverside as a model species for identifying estuarine environments that have reduced embryo viability due to contamination. My hypothesis is that Atlantic silverside embryos collected from contaminated estuaries, such as Murrells Inlet, will have a lower percent viability compared to those collected in protected estuaries such as the North Inlet estuary.

NERR emergent vegetation bio-monitoring: Effects of sea level on the spatial dynamics of salt marsh vegetation communities in the North Inlet estuary

Investigators: Tracy Buck, Erik M. Smith, Julie L. Krask, and Baker Stevens

North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

As part of a National Estuarine Research Reserve (NERR) system-wide initiative, the North Inlet–Winyah Bay NERR is monitoring salt-marsh emergent vegetation with the aim of quantifying variability in salt marsh macrophyte community spatial structure (species composition, relative abundance, and biomass) along elevation gradients, from creek bank to upland edge, in response to changes in tidal height and flooding frequency due to sea level rise. Long-term monitoring is conducted in accordance to established NERRS protocols using a stratified sampling approach of fixed transects and repeated measures within permanent sample plots. This consists of two marsh segments with three fixed transects and 20 sampling plots per transect. Surface Elevation Tables (SETs) have also been established adjacent to the lower and higher elevations of the creek-bank to forest-edge transects in each marsh region to determine changes in marsh surface elevation associated with long-term changes in and vegetation and tidal dynamics. Sampling within each permanent plot includes: percent cover for each species or cover category; species' shoot/stem density; species' maximum canopy height; species' aboveground biomass by non-destructive sampling techniques; water table height at low tide; porewater salinity, and nutrient and sulfide concentrations. Soil organic content and bulk density adjacent to each plot were determined in 2008. Elevation data (mm scale vertical resolution) for each plot is determined at biannual intervals to allow for the calculation of duration and frequency of tidal inundation at each plot.

Shorebird monitoring in the North Inlet estuary

Investigators: Jennifer Plunket and citizen scientist volunteers Wendy Allen and Emma Boyer

North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

Shorebirds (Aves: Charadriiformes), are a diverse group of birds including plovers, sandpipers, curlews and oystercatchers. Of the more than 50 different species that occur in North America, more than half are considered a species of concern or “highly imperiled” due to declining numbers. A shorebird monitoring effort was initiated in the spring of 2016 to assess populations of shorebirds in the North Inlet estuary, primarily during migration periods, March - June and July - October. Shorebird surveys in the North Inlet estuary are conducted biweekly during these periods near high tide by boat and land and through the use of wildlife cameras. Species are identified and counted at sites on Hobcaw Beach, North Island, North Jones Creek, Bosun's Point, and along Clambank Causeway. Survey data is entered into a database using protocols established by the International Shorebird Survey (ISS) administered by the Manomet Center for Conservation Services. Color-marked individuals are also noted. This project will help establish baseline information on the species and numbers of shorebirds utilizing the North Inlet estuary during periods of migration and will help to identify key habitat areas within the reserve. It will also feed into the larger ISS database that includes information from about 1,200 locations in North America that is contributing to a better understanding of shorebird population numbers, key stopover locations, migratory routes, and other aspects of shorebird life histories

Characterization of oyster cement

Investigators: Jonathan Wilker¹ and Paul D. Kenny²

1 - Department of Chemistry, Purdue University (IN)

2 - Baruch Marine Field Laboratory, University of South Carolina

Marine species such as mussels, barnacles, and oysters produce adhesive and cement materials for affixing themselves to surfaces. The strong bonding, wet adhesion capabilities, and biological origin of these materials indicate promise for developing new biomedical materials such as surgical glues and dental cements. In an effort to develop such applications, we are beginning by characterizing adhesive materials produced by marine organisms. Prior studies have determined some of the key chemical reactions and bonding motifs used by mussels for production of their adhesive. For the current project, our main objective is to characterize the chemistry within the cement of the Eastern or Atlantic oyster *Crassostrea virginica*. Oysters are collected near the Baruch Marine Field Laboratory and then grown in laboratory aquaria. Chemical methodologies are used to analyze the cement, including wet chemistry and spectroscopic techniques. Insights gained will provide both fundamental understanding of how a marine biological material functions as well as providing insights for the design of new biomedical adhesives.

The effects of inhibited carbonic anhydrase on the phytoplankton communities in coastal waters

Investigators: Eilea Knotts¹ and James L. Pinckney²

1 - Department of Biological Sciences, University of South Carolina

2 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

Carbon concentrating mechanisms (CCMs) are used by phytoplankton in order to concentrate dissolved inorganic carbon (DIC) within their cells. These adaptations were evolved to enhance uptake of DIC at present surface water concentrations. However, mechanisms, such as the carbonic anhydrase enzyme (CA), are active, energy-consuming processes that may become redundant in the future due to increased concentrations of CO₂ in surface waters. In order to gain a better understanding of the carbonic anhydrase enzyme, recent studies have investigated the CA enzyme through the use of inhibitors: acetazolamide (AZ) and ethoxzolamide (EZ). Most of our knowledge is based on individual cultures or oligotrophic water samples. However, there are few studies that look at the mechanism's effects on estuarine phytoplankton communities and none have measured the *in situ* effects on community composition. Using bioassays of natural phytoplankton communities, our research will explore how community composition is altered when the competitive advantage of the CA enzyme is reduced. These changes will be monitored with measurements of chlorophyll *a* fluorescence, cell abundances, microscopy, photopigments, nutrients, and the inhibitors: AZ and EZ. This study will provide a better understanding of how the CA enzyme impacts the composition of phytoplankton communities in coastal waters.

Evaluating intertidal oyster reef restoration success

Investigators: Keith Walters, Thomas Funk, and students

Department of Marine Science, Coastal Carolina University (SC)

A series of oyster reefs, shell-filled mesh bags, were created within salt marsh-estuaries (Hog, Murrells, and North Inlets) and swash tidal creeks (Whitepoint, Singleton, and Withers) to evaluate the success of reef restoration efforts. In the North Inlet estuary, reefs were created within Bly Creek in June 2014 and, along with coincident natural reefs, have been the focus of graduate and undergraduate researchers and CCU classes (e.g., Marine Ecology). The following data were or continue to be collected to evaluate constructed reef development and assess the ability of inlet and swash constructed and natural reefs to attract and support important fishery species: (1) yearly oyster spat recruitment; (2) numbers, sizes, and distribution of oysters on constructed and natural reefs; (3) numbers and species richness of resident and transient nekton associated with reefs; (4) short-term predation on reef-resident bivalve and decapod fauna. Although oyster recruitment onto reefs is similar, preliminary results suggest inlet oysters survive better as juveniles and live longer as adults compared to swash creek populations. Nekton seasonally captured during high tides within baited minnow traps, pull traps, and gill nets indicate constructed reefs are colonized quickly. Survival of tethered bivalve and decapod individuals within mudflat and constructed and natural reefs documented initial differences in structural complexity and the role of complexity in the survival of reef-resident taxa. The ongoing studies are providing valuable experiences for students, identifying demographic differences that influence reef development, and demonstrating the value of reef restoration efforts to intertidal resident and transient taxa.

A collaborative science program for the National Estuarine Research Reserve System (NERRS): Working with end users throughout the applied research process

Investigators: Dwayne E. Porter^{1,2}, Melissa Ide³, and Jeremy Cothran¹

1 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

2 - Arnold School of Public Health, University of South Carolina

3 - Centralized Data Management Office, National Estuarine Research Reserve, University of South Carolina

The NOAA NERRS Science Collaborative (NSC) supports integrative environmental and social research for improved community decision making. The NERRS Centralized Data Management Office (CDMO), housed at the Baruch Marine Field Laboratory, is the lead for University of South Carolina's (USC) involvement in the establishment and administration of the NOAA NERRS Science Collaborative (NSC). The NSC is led by researchers at the University of Michigan, Stanford University, and USC. The CDMO role is the transfer of key knowledge and lessons learned to others, potentially benefiting NERRS as well as local, state and federal coastal management decision makers and educators; and delivery of highly credible, valid and relevant scientific results and data that are both timely and universally accessible.

Long-term measurements of production and physiological ecology of *Spartina alterniflora*

Investigators: James T. Morris¹ and Karen Sundberg²

1 - Department of Biological Sciences, University of South Carolina

2 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

Salt marsh grass (*Spartina alterniflora*) dominates the intertidal marsh in North Inlet estuary. Regular measurements of grass density and height allow for estimates of growth and primary production rates in both control and fertilized plots. Abiotic conditions that are measured include pore water salinity, phosphate, ammonium, sulfide, and iron concentrations to provide insights into factors that affect production. Large monthly and interannual variations in the amount of organic material produced by the cordgrass are related to such factors as sea level and precipitation patterns. This time series was initiated in 1986.

Co-evolution of complex traits associated with key innovation: Armor and vision in the snapping shrimp (*Alpheus heterochaelis*)

Investigators: Alexandra C. N. Kingston, Daniel I. Speiser, and Rebecca Lucia

Department of Biological Sciences, University of South Carolina

Snapping shrimp are a morphologically diverse and speciose group of crustaceans in which the evolution of their specialized snapping claw is accompanied by the evolution of protective armor. This armor, called the orbital hood, conceals the eyes of snapping shrimp to protect them from the explosive collapse of cavitation bubbles produced by their snapping claws. To study the co-evolution of these traits, we examined the optical properties of the head-protecting armor, the morphology and physiology of the visual system, and the visually-influenced behaviors of *Alpheus heterochaelis*. We found that the orbital hood of *A. heterochaelis* is highly transparent, transmitting 80-90% of incident light. Electroretinography shows that *A. heterochaelis* responds maximally to 500 nm light and demonstrates a flicker fusion frequency of > 40 Hz. Microspectrophotometry reveals the presence of two middle-wavelength sensitive visual pigments, $\lambda_{max} = 500$ and 519 nm. Optomotor assays show that snapping shrimp demonstrate spatial vision: animals will follow moving stripes with angular widths as narrow as 4°. These results indicate that *A. heterochaelis* has a functional visual system that provides spatial vision despite the presence of the orbital hood. We will continue investigating the relationships between the weapon, armor, and visual system by exploring pressure generation by the snapping claw and the effect of snapping on orbital hoods during agonistic interactions. Studying protective armor in an animal with such a powerful weapon provides opportunities to examine relationships between these and other complex traits, and the evolutionary trade-offs that result from the evolution of specialized morphological traits.

Long-term monitoring of grass shrimp as a bioindicator of non-point source runoff in South Carolina watersheds

Investigators: Peter Key, James Daugomah, and Blaine West

Center for Coastal Environmental Health and Biomolecular Research, National Oceanic and Atmospheric Administration (SC)

Long-term ecological monitoring is important to developing fundamental understandings of both biogenic and anthropogenic effects on ecosystem health. Long-term monitoring may provide great insight into natural factors such as disease, pests, and weather (e.g., global climate change, drought, floods, and increased intensity of tropical storms and hurricanes), which may affect populations throughout a geographical region. In addition to population perturbations caused by natural stressors, is the complexity of differentiating anthropogenic effects of chemical and biological contaminants in aquatic ecosystems from natural background effects. There is a clear need to develop accurate ecological forecasts using long-term ecological data sets. Long-term ecological monitoring data thus can be used not only to ascertain effects of natural and anthropogenic stressors, but also when properly used in conjunction with GIS and advanced modeling techniques may enhance predictive capabilities. The grass shrimp (*Palaemonetes pugio*) is the dominant motile macrobenthic invertebrate in tidal creek systems of the southeastern United States and is an important prey item for higher trophic levels. The Oyster Landing site within the North Inlet estuary is maintained as a long-term reference site for comparison to estuarine sites with other land uses.

The National Estuarine Research Reserve System Centralized Data Management Office

Investigators: Dwayne E. Porter^{1,2}, Melissa Ide³, Jennifer Kesse³, Amber Knowles³, Brooks Folk³, Lee Shutt³, Dan Ramage², and William H. Jefferson¹

1 - Belle W. Baruch Institute for Marine and Coastal Sciences, University of South Carolina

2 - Arnold School of Public Health, University of South Carolina

3 - Centralized Data Management Office, National Estuarine Research Reserve, University of South Carolina

NOAA's National Estuarine Research Reserve System (NERRS) acknowledges the importance of both long-term environmental monitoring programs and data and information dissemination through the support of the NERRS System-Wide Monitoring Program (SWMP). The goal of the SWMP is to "identify and track short-term variability and long-term changes in the integrity and biodiversity of representative estuarine ecosystems and coastal watersheds for the purpose of contributing to effective national, regional and site specific coastal zone management." This comprehensive program consists of three phased components: estuarine water quality monitoring (phase I), biodiversity monitoring (phase II), and land-use and habitat change analysis (phase III). The Centralized Data Management Office (CDMO) was established in support of the System-Wide Monitoring Program involving 29 sites around the US and Puerto Rico. The purpose of the CDMO, housed at the North Inlet-Winyah Bay NERR, is the management of the infrastructure and data protocol to support the assimilation and exchange of data, metadata, and information within the framework of NERRS sites, coastal zone management programs, and other education, monitoring and research programs.

Effect of wrack accumulation on salt marsh vegetation, Baruch Institute, Georgetown County, South Carolina

Investigators: Richard Stalter¹, A. Jung¹, A. Starosta¹, John Baden², and M. D. Byer³

1 - Department of Biological Sciences, St. John's University (NY)

2 - US Army Corps of Engineers, NC, Retired

3 - Gateway National Recreation Area, National Park Service (NY)

In March 2004, four arrays in different types of salt marsh vegetation were covered with 15-20 cm of wrack secured in place, in an attempt to duplicate the natural deposition of wrack on the marsh by tides and storms, and to quantify and extend anecdotal observations and the results of previous studies. A control plot in each array was left uncovered; another plot was covered with only 2-3 cm. The wrack was removed from one plot in each array at one, two, four and seven month intervals. One month of wrack coverage appeared to have little effect on either density or standing crop, recorded seven months after initial covering, of the principal marsh species. Above ground parts of these species, with the exception of *Spartina patens* and *Borrichia frutescens*, appeared to be killed or extremely inhibited after two months, but not as much as the 15-20 cm wrack mat treatment. We continue to assess survival of wrack impacted plants and monitor recruitment and growth in specific wrack impacted zones.

Estimating above ground biomass utilizing Planet multispectral satellite data

Investigators: Gwen Miller¹ and James T. Morris²

1 - School of the Earth, Ocean, and Environment, University of South Carolina

2 - Department of Biological Sciences, University of South Carolina

Conducting field work to analyze landscape scale above ground biomass within the North Inlet-Winyah Bay National Estuarine Research Reserve (North Inlet) is infeasible due to the extent of the salt marsh and access difficulty. Estimates of above ground biomass is important for better understanding marsh productivity and modeling how a salt marsh may respond with sea-level rise. Furthermore, a landscape scale map of above ground biomass within North Inlet will highlight regions within the marsh with higher or lower biomass. Publicly available satellite data such as data available through Landsat, have a large spatial resolution (30 meters), which smooths out spatial variability. A private company, Planet, offers us free 3-meter resolution multispectral through Planet's education and research program. We will collect above ground biomass samples at various locations within the North Inlet and use Planet's spectral data to create a model between above ground biomass and satellite data. Using the calibrated model, we will develop a map of above ground biomass within the entire salt marsh dominated by *Spartina alterniflora*.

Hard clam (*Mercenaria mercenaria*) population dynamics in North Inlet estuary tidal creeks

Investigator: Juliana M. Harding

Department of Marine Science, Coastal Carolina University (SC)

Hard clam (*Mercenaria mercenaria*) populations play an ecological and structural role within tidal creek habitats. The population biology and dynamics of hard clams are being quantitatively examined in North Inlet estuary tidal creeks. Hard clam age structure, growth rates and sex ratios continue to be evaluated seasonally and combined with measurements of environmental variables to describe clam population dynamics in tidal creeks and their effects on habitat structure within the creeks over multi-year time scales.

Physical characteristics of estuarine waters: Long-term monitoring in the North Inlet and Winyah Bay estuaries

Investigators: Erik M. Smith, Baker Stevens, Julie L. Krask, and Tracy Buck

North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

As part of the NERR System-Wide Monitoring Program, the physical characteristics of the water in four tidal creeks of the North Inlet–Winyah Bay NERR have been monitored using YSI data loggers since 1994. A new, fifth site in the mainstem of Winyah Bay was added in 2016. These data loggers are deployed at 0.5 m above the sediment surface and record water depth, temperature, salinity, pH, dissolved oxygen, and turbidity at 15 min intervals throughout the year. The site in Winyah Bay has data loggers deployed in both surface and bottom waters to account for the vertical stratification that exists in this location. The instruments are calibrated and deployed according to strict NERRS protocols. The consistent, long-term collection of this physical data allows for the characterization of short-term variability and long-term change in North Inlet waters, and provides base-line data critical for various studies of biological and physical processes in the North Inlet estuary. Data, along with detailed metadata, are sent to the NERRS Centralized Data Management Office (CDMO) for quality assurance and quality control. Data can be accessed via the CDMO website (<http://cdmo.baruch.sc.edu/>).

Mosquitoes of the Carolinas

Investigator: Brian Byrd

School of Health Sciences, Western Carolina University (NC)

The overall project seeks to monitor the presence of native and invasive mosquito species in NC and SC. Particular focus is on the spread of traditionally southern species that may be moving north and the presence of invasive mosquito species. Hobcaw Barony represents an ideal collection and monitoring site with more than 20 known mosquito species obtained by the principal investigator during previous collections. The mixed land use and land classifications at Hobcaw provide reliable locations for the collection of salt marsh, permanent and semi-permanent freshwater wetland mosquito species. Mosquito specimens will be obtained by larval sampling, aspiration, and light traps.

Analyzing the seasonal change of heavy metal sediment content in Winyah Bay estuary tidal marshes

Investigators: Isabel Fangman and Jennifer Hawk

Department of Biology, Chemistry, and Physics, Converse College (SC)

Wetland areas are an extremely vital part of the environment and serve a variety of functions including filtering out pollutants in the water system and providing food sources for humans. Heavy metals can enter the food chain through the wetlands and accumulate in toxic levels in the plants and animals that live there. To help monitor the human influenced stress-levels of heavy metals in the environment, the Environmental Protection Agency (EPA) established the Heavy Metal Index. The goal of this study is to measure the heavy metal content in sediment in tidal marshes of the Winyah Bay estuary on the Hobcaw Barony property in Georgetown, SC. This study will help establish a baseline for the heavy metal content in this area. Sediment samples will be collected from the marshlands and analyzed using Atomic Absorption Spectrometry. The levels of chromium (Cr), lead (Pb), iron (Fe), nickel (Ni), copper (Cu), and cobalt (Co) will be studied over the course of three seasons, specifically spring, summer, and fall, to determine if and how those levels change over the course of the year.

Examination of potential juvenile tarpon (*Megalops atlanticus*) overwintering habitat in the North Inlet estuary

Investigators: Matthew E. Kimball, Dennis M. Allen, and volunteers
Baruch Marine Field Laboratory, University of South Carolina

Juvenile tarpon occur in small, shallow, and often tidally restricted environments in and around estuaries (e.g., ditches, pools or ponds, and impoundments) during summer and fall. Based on laboratory low temperature tolerance experiments, juvenile tarpon that do not emigrate in late fall likely do not survive the winter in most estuarine habitats where temperatures drop below their lethal limit (~10-13° C). However, tarpon have been observed during winter months in a thermally stratified pond with little tidal influence, as well as artificially warmed inshore habitats (e.g., discharge canal from power plant). To examine the overwinter potential of thermally unique estuarine habitats, we have monitored the survival and growth of juvenile tarpon in a pond along the upland edge of the North Inlet estuary since 2016. Juvenile tarpon (~100 - 500 mm SL) have been consistently observed in this pond and preliminary temperature data suggests that it may provide suitable overwinter habitat. Throughout the year, juvenile tarpon are collected using hook-and-line, tagged with a 12-mm Passive Integrated Transponder (PIT) tag, and released back into the pond. Both pond surface and bottom water temperatures are measured continuously during winter months (December through March). Combined, information on tarpon habitat use garnered through observations of tagged individuals along with winter water temperature records will provide a better understanding of tarpon survival in potential overwinter habitat in this region.

Development of models for phytoplankton-nutrient responses in support of numeric nutrient criteria for estuarine water quality

Investigators: James L. Pinckney¹, Erik M. Smith², and Krystyn Kibler³

1 - Belle W. Baruch Institute of Coastal and Marine Science, University of South Carolina

2 - North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

3 - School of the Earth, Ocean, and Environment, University of South Carolina

The primary objective of the proposed research is to develop and test a family of empirical mathematical models to quantify the responses of the total phytoplankton community as well as phytoplankton groups to increases or reductions in total nitrogen (N) loading. Models will be constructed for a range of N loading scenarios under both high and low light exposure conditions. These models will be invaluable for developing and validating numeric nutrient criteria for Winyah Bay and the North Inlet estuary and provide a “proof of concept” for determining criteria in other estuarine systems. Phytoplankton-nutrient response curves will be constructed using natural phytoplankton communities collected in the North Inlet estuary and Winyah Bay, South Carolina (SC). Surface water (0.5 m depth) will be obtained seasonally over 2 years from each location. The N treatment for the bioassays will consist of a range of concentrations (1 - 100 $\mu\text{mol N l}^{-1}$ in increments of 10 $\mu\text{mol N l}^{-1}$) and will be composed of an equimolar mixture of NO_3^- , NH_4^+ , and urea ($\text{CO}(\text{NH}_2)_2$) to simulate the types of N compounds likely available in the estuary. Data from the bioassays will be used to derive an empirical numerical relationship between N loading and phytoplankton community biomass (as chl *a*) responses. The empirical models we will develop can be directly used by the SC Department of Health and Environmental Control to evaluate numeric nutrient criteria for these systems under a variety of N addition/reduction scenarios.

Benthic foraminifera as indicators of ecosystem health in the North Inlet estuary

Investigators: Kelly Gibson and Sarah Shealy

Department of Biology and Geology, University of South Carolina Aiken

The distribution of benthic foraminifera (single-celled benthic microorganisms that secrete a calcite shell) has been directly tied to environmental parameters such as salinity, nutrient, and dissolved oxygen content. We intend to establish relationships between environmental parameters (i.e., salinity, sediment grain size and organic matter content) in the North Inlet estuary and faunal assemblages of benthic foraminifera by collecting surface sediment samples for faunal and geochemical analyses. The North Inlet estuary is relatively pristine and will serve as a baseline with which to compare conditions in estuarine environments with more severe anthropogenic influences/impacts in future studies, both spatially and temporally.

Satellite observations of marsh dieback events and potential environmental influences along coastal South Carolina, 2000-2017

Investigators: Cuizhen (Susan) Wang and Jean T. Ellis
Department of Geography, University of South Carolina

Using the North Inlet estuary salt marsh as one of the experimental sites, this study aims to identify the marsh dieback events and recovery pathways in South Carolina (SC) via a series of National Aeronautics and Space Administration (NASA) satellite observations since 2000. SC has over 340,000 acres of salt marshes that support 96% of the state's commercial seafood and shellfish catch. Since 2000, studies have reported multiple dieback events in the U.S. Southeast. Investigations of marsh dieback in SC, however, have been limited. The most pristine along the east coast, the SC marshes and dieback-recovery pathways deserve critical research. Satellite images provide a long-term, continuous way to identify the thinning and dead marsh patches. Supported by the NASA EPSCoR program, this study will address the following objectives to: 1) classify marsh dieback patches through Landsat (TM, ETM+, OLI) image series; and 2) identify initial indicators of stressors by exploring the environmental anomalies during the events from NASA satellite products (MODIS; TRMM-GPM; Aquarius-SMAP). The expected outcomes of this project include an inventory of dieback events and potential stressors in SC. When future funding opportunities are available, these results will help us explore how these environmental stressors impact marsh healthiness, and how marsh habitats sustain under natural and anthropogenic impacts along coastal SC.

American alligator (*Alligator mississippiensis*) movement ecology in coastal South Carolina

Thomas R. Rainwater¹, Abigail J. Lawson², and Patrick G.R. Jodice³

1 - Baruch Institute of Coastal Ecology and Forest Science, Clemson University (SC)

2 - Department of Forestry and Environmental Conservation, Clemson University (SC)

3 - U.S. Geological Survey South Carolina Cooperative Fish and Wildlife Research Unit, Department of Forestry and Environmental Conservation, Clemson University (SC)

Effective wildlife management necessitates a working knowledge of demography and habitat requirements. In South Carolina (SC), the American alligator (*Alligator mississippiensis*) is managed as a renewable natural resource, through public and private lands harvest programs. To date, SC's monitoring program for alligators has primarily focused on an estimation of the total number of animals in the state-wide population but has not incorporated efforts to identify underlying mechanisms that may drive variation in realized or observed abundance patterns (e.g., habitat selection, associated movement patterns). In 2015, we initiated a movement ecology study of adult male alligators in coastal SC and deployed GPS transmitters on animals in the Santee Delta and ACE Basin regions of the state. During our tracking efforts, we observed some alligators move significant distances over short periods. One animal captured and tagged at the Tom Yawkey Wildlife Center routinely crossed Winyah Bay and spent considerable time on the southern end of the Hobcaw Barony (HB) property, highlighting both extensive movement as well as the potential importance of habitats in the HB/North Inlet estuary area for coastal alligators. In 2017, we tagged an adult male alligator at HB and have tracked its movements for the last year. Results indicate this animal uses multiple freshwater and estuarine habitats on HB, complementing data from other alligators tracked in coastal SC. We now aim to investigate how alligators select and utilize these different habitat types (e.g., foraging, breeding, nesting, overwintering) using a combination of GPS telemetry, trail cameras, and stable isotope analysis.

Painted bunting (*Passerina ciris*) breeding survey

Investigators: Jennifer Plunket¹ and citizen scientist volunteers Wendy Allen, Bill Brabson, and Marlene Konsek

1 - North Inlet-Winyah Bay National Estuarine Research Reserve, University of South Carolina

The project goal is to estimate of the number of painted buntings utilizing the edges of North Inlet estuary marshes as nesting habitat during the breeding season. A point count method is utilized that involves 5 minutes of listening for calling male buntings at fixed intervals along a prescribed route. Summer 2019 marked the second year of this survey. The plan is to continue this breeding bird survey of painted buntings each year so that changes in numbers can be detected for this species of high concern in South Carolina. The point count methodology is consistent with North American Breeding Bird Surveys and a population assessment of painted buntings conducted in the southeast, 2007-2009, thus allowing for comparisons with other studies.

Determining age structure of juvenile tarpon (*Megalops atlanticus*) in nursery habitats in South Carolina estuaries

Investigators: Garrett Elmo¹, Matthew E. Kimball², and Derek P. Crane¹

1 - Department of Biology, Coastal Carolina University (SC)

2 - Baruch Marine Field Laboratory, University of South Carolina

Recruitment of early life stage tarpon to coastal nursery areas in this region typically peaks during the summer months, and tarpon rear in a variety of marsh habitats that range widely in water quality (e.g., salinity, dissolved oxygen) and physical (e.g., substrate, depth) characteristics. Size differences between juvenile Tarpon in natural marsh habitats compared to marsh impoundments (about 2.5 times bigger on average) were recently (2016) documented in the North Inlet-Winyah Bay estuarine system. These size differences occurred despite larval tarpon recruiting to both habitats around the same time (June). While these size differences may be due to factors such as habitat quality and varying growth rates, discerning the age structure of early life stage tarpon populations in these habitats is a critical first step for understanding recruitment and habitat use patterns, and ultimately determining the underlying mechanisms for such observed size differences. We will estimate the age of juvenile tarpon from natural marsh habitats and marsh impoundments using extracted sagittal otoliths. Prior to processing otoliths for age estimation, dried otoliths will be weighed, as earlier work indicated that otolith weight was positively correlated with age for juvenile tarpon. After weighing, otoliths will be mounted in epoxy and cut into transverse sections containing the core. Sections will then be mounted to a glass slide and polished until daily growth increments are visible. Otoliths will be viewed using a zoom stereo microscope and associated software, and daily increments will be counted and used to determine potential age differences between habitats.

Relating nonpoint source BOD loading to land-use and stormwater management practices in coastal South Carolina

Investigators: Erik M. Smith¹, Claudia Benitez-Nelson², and Kelly McCabe²

1 - North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

2 - School of the Earth, Ocean, and Environment, University of South Carolina

The presence of a sufficient concentration of dissolved oxygen (DO) is a fundamental requirement necessary to sustain aquatic life. The occurrence of low DO is the number one cause of impairment with respect to the ability of waters to support aquatic life use in South Carolina's coastal zone, representing 48% of all EPA Section §303(d) listings for Aquatic Life Use impairment in the eight coastal counties. Maintaining sufficient DO requires managing anthropogenic discharges of oxygen demanding substances. The operational measure of oxygen demanding substances is the determination of five-day Biochemical Oxygen Demand (BOD). While BOD loading from point sources is relatively straight-forward, the role of nonpoint sources in contributing BOD loading to coastal waters remains a significant information gap. The overall goal of this research is thus to develop a predictive understanding of how land-use and specific stormwater management conveyances and structural control structures impact variations in the magnitude of nonpoint source BOD. This will be accomplished by analyzing BOD on samples collected from 1) first-order catchments that represent the range of land cover and land use found in this region, from largely natural (including both forested uplands and wetlands) to high density urban development; and 2) the outfalls of common stormwater management practices (detention ponds, constructed wetlands, etc.). Water samples analyzed for BOD will also be subject to a suite of carbon, nitrogen and phosphorus analyses. Anticipated results will contribute to both improved stormwater management and Total Maximum Daily Load (TMDL) development for coastal waters impacted by low DO.

3-D subsurface modeling of a clastic paleo-shoreface deposit

Investigator: Patrick Duff

School of Earth, Ocean, and Environment, University of South Carolina

Close examination of modern shallow marine deposits can also help us better understand similar ancient deposits, including their internal structure and relationship to other passive margin deposits. A series of Ground Penetrating Radar (GPR) lines, along with vibrocores, and a UAS drone survey will be acquired near the Baruch Marine Field Laboratory on Hobcaw Barony, as one field site within the coastal plain of South Carolina to investigate the internal structure and stratigraphy of a variety of a recent shoreface deposit, as well as to serve as an experiment in the application of GPR to sedimentological and stratigraphic research.

Satellite observations of marsh dieback events and spatial assessment of environmental influences

Investigators: Huixan Li and Cuizhen (Susan) Wang
Department of Geography, University of South Carolina

The primary goal of this research is to detect the marsh dieback events from the long-term satellite observations (since 2000) and to examine the potential influencing factors by investigating the spatial and temporal relationships between marsh diebacks and satellite-observed environmental anomalies. To achieve this goal, the following objectives will be addressed: 1) Map the current distributions of primary marsh species from recent Landsat images in the two National Estuarine Research Reserves; 2) Identify the dieback patches in marshes from Landsat (TM, ETM+, OLI) image series in 2000-present that experienced apparent greenness reduction, indicating marsh dieback events; 3) Explore the relationships between these dieback events and environmental anomalies from National Aeronautic and Space Administration (NASA) satellite products (MODIS; TRMM-GPM; Aquarius-SMAP) and in-situ water quality data records, which are primary indicators of environmental influences; 4) Perform comparison analysis of marsh dieback events between the two study sites (natural vs. land development) to examine the anthropogenic impacts on marsh dynamics on SC coasts.

Weather and climate measurements: Long-term monitoring at Oyster Landing pier

Investigators: Erik M. Smith, Baker Stevens, Julie L. Krask, and Tracy Buck
North Inlet–Winyah Bay National Estuarine Research Reserve, University of South Carolina

As part of the North Inlet–Winyah Bay National Estuarine Research Reserve (NERR), a fully functional meteorological station (National Weather Service installation) is located on the Oyster Landing pier in the North Inlet estuary. Wind speed and direction, air temperature, humidity, barometric pressure, solar radiation, and precipitation are recorded at 15-minute intervals. Data are telemetered via the NOAA GOES satellite system to the NERR Central Data Management Office, and made available in near real time (<http://cdmo.baruch.sc.edu>). For most parameters, records have been collected for more than 20 years. Long-term, continuous weather records provide data for determining the effects of climatology on the various biological and physical processes being studied in the North Inlet estuary.

Diet of terrestrial-mammalian carnivores that have access to marine-aquatic resources compared to terrestrial upland habitats in coastal South Carolina

Investigators: Amanda L. Wilson¹, David Jachowski¹, Ernie Wiggers², Michael Childress³, Patrick Gerard⁴, and Lisette Waits⁵

1 - Department of Forestry and Environmental Conservation, Clemson University (SC)

2 - Nemours Wildlife Foundation (SC)

3 - Department of Biological Sciences, Clemson University (SC)

4 - Department of Mathematical Sciences, Clemson University (SC)

5 - Department of Fish and Wildlife Sciences, University of Idaho

Coastal ecotones, where marine aquatic and terrestrial habitats interact, have been shown to be productive systems where allochthonous input subsidize terrestrial mammalian predators. Research involving the diets of predators along the Atlantic coast is sparse, yet important because it allows for a fuller understanding of community dynamics, and function and position of animals within a food web. The objectives of this study are to compare diets of predators that have access to coastal ecotones to upland areas in order to (1) assess how the diet of each predator using coastal areas (e.g. impoundments) is influenced by allochthonous inputs, (2) document any seasonal variation in its use, and (3) compare dietary breadth and overlap among predators that could inform community ecology of sympatric predators' dynamics and structure. By conducting scat surveys across the coastal plain of South Carolina, this study will compare the diets of those predators, via scat, using microscopic and macroscopic identification of prey items from reference collections. Further, this study will use molecular genetic techniques to confirm predator species identification since rates of misclassification using field morphometric identification of scat is high. This research will contribute to a gap of basic ecological knowledge of how predators utilize these coastal ecotones and will inform land managers and property owners of how predators interact with those species for which properties are predominantly managed.

Validation of back-calculation methods for juvenile tarpon (*Megalops atlanticus*)

Investigators: Garrett Elmo¹, Derek Crane¹, Matthew Kimball²

1 - Department of Biology, Coastal Carolina University (SC)

2 - Baruch Marine Field Laboratory, University of South Carolina

Biologists use growth as a metric to measure habitat quality under the assumption that good growth equates to good habitat for a species and their prey. Therefore, by understanding growth and the factors affecting growth, biologists can determine which habitat types are important to protect and restore for conservation of early life stages of many fish species. Size differences between juvenile tarpon in natural marsh pools in the North Inlet estuary and marsh impoundments in the Yawkey Wildlife Preserve were recently documented. Fish size and growth during the juvenile life stage can be affected by a variety of biological and environmental factors including: age of fish, diet and prey quantity/quality, salinity, temperature, and competition. However, drivers of differences in juvenile tarpon size observed in small, shallow, and tidally-restricted habitats in natural and managed marshes in this region are unknown. Growth of fishes is commonly estimated by (1) estimating mean length-at-age, after assigning ages based on counting annuli in calcified structures (e.g., bones, otoliths, scales, fin rays; synonymous with counting rings in a cross-section of a tree), (2) using measurements between annuli in calcified structures to back-calculate length-at-age, or (3) using capture-recapture methods. Capture-recapture methods typically have lower sample sizes because of the difficulty of recapturing marked fish from an open system. Recapture rates of juvenile tarpon are relatively low, therefore methods 1 and 2 are better suited for estimating juvenile tarpon growth. Both methods assume ages can be estimated accurately from calcified structures and back-calculation assumes calcified structure growth and fish growth have a linear relationship. To date, no studies have examined the validity of back-calculation methods for tarpon, and thus we propose to evaluate the accuracy of back-calculation methods here.

Enhancing bivalve aquaculture through species improvement and diversification

Investigators: Ximing Guo¹, Daphne Munroe¹, Lisa Calvo¹, David Bushek¹, Michael DeLuca², and Pete Rowe³

1 - Haskin Shellfish Research Laboratory, Rutgers University (NJ)

2 - Aquaculture Innovation Center, Rutgers University (NJ)

3 - New Jersey Sea Grant Consortium

Bivalve aquaculture is one of the most important aquaculture industries in the US and globally. It is environmentally friendly, family or community-based, and important to the socioeconomic recovery of coastal areas suffering from declining fisheries. Aquaculture has been growing steadily; however, major impediments - diseases, unpredictable mortality and a lack of crop diversity - limit the long-term growth and sustainable future of the bivalve aquaculture industry. This project seeks to diversify the selected varieties of oysters available to the Atlantic coast aquaculture industry. Bivalve aquaculture in the US is dominated by three species: the eastern oyster (*Crassostrea virginica*) and hard clam (*Mercenaria mercenaria*) along the Atlantic and Gulf coasts, and the Pacific oyster (*Crassostrea gigas*) along the Pacific coast. US aquaculture production of the three species was valued at 94, 46 and 63 million dollars, respectively, in 2015, which represent 133%, 1% and 13% increases compared with that in 2005. The eastern oyster and hard clam support major aquaculture industries along the Atlantic coast. Both species are threatened by diseases but existing disease tolerant strains have been selected to perform in low to moderate salinity limiting their performance in other areas. Another major impediment is the lack of crop diversity. Diversifying the species grown is especially important for aquaculture growth and resilience under the threat of diseases and unpredictable mortality. This project selected oysters from North Inlet estuary, SC as a source of high salinity, warm temperature, intertidally adapted broodstock to improve strain diversity for aquaculture.

Chemical analysis of soil samples to detect events in a linear model

Investigators: Gabrielle Johnson, Ashley Davis, Samantha McClain, and Students Engaged in Aquatic Sciences members

School of the Earth, Ocean, and Environment, University of South Carolina

This project aims to measure carbon sequestration and nutrient accumulation in marsh sediments, and to introduce new members of the undergraduate program Students Engaged in Aquatic Sciences (SEAS) to sediment geochemistry. A transect will be taken with three cores on the northwest and three cores on the southeast sides of Bly Creek in the North Inlet estuary to get a spatial distribution ranging from forest towards the ocean. Collected sediment cores will be analyzed for carbon and nitrogen to determine productivity and lead-210 to determine the age.

Associations between temperature and larval recruitment of a non-native crab (*Petrolisthes armatus*) near its northern range limit

Investigators: Teresa Popp and Dara Wilber
Grice Marine Lab, College of Charleston (SC)

The non-native green porcelain crab, *Petrolisthes armatus*, was first seen on intertidal oyster reefs in South Carolina in 1995. Since then, its range has expanded north in what appears to be a series of surges and setbacks associated with cold winters that may cause mortality. *P. armatus* reproduces year-round in the native range but is limited to warmer months in the novel range. Variation in the timing and duration of larval recruitment are not known for the introduced range and these factors may affect further range expansion. Our research examined whether *P. armatus* larval recruitment is related to temperature variation, in particular, the severity of winter temperatures. Biweekly larval tows collected in the North Inlet–Winyah Bay National Estuarine Research Reserve since 1981 provide a valuable resource for assessing larval recruitment trends. We analyzed samples collected from 1994 to 2018 to determine whether the timing and duration of larval ingress varied over time and related to temperature variation. *Petrolisthes armatus* larvae reached the region in 1995 and once established, has been present ever since. Results also show there is considerable variability in when recruits are first observed (coefficient of variation 29%) and that earlier annual recruitment was positively correlated with a longer duration of recruitment ($r = 0.64$, $p = 0.001$). The timing of larval recruitment was also related to the severity of the preceding winter, where later larval recruitment occurred in years with several consecutive days of cold temperatures (7-day minimum temperature; $r = 0.44$, $p = 0.030$). These results support the theory that cold winter temperatures affect overwintering populations and their ability to reproduce in the early spring. However, the frequency of severe winters in the study area is decreasing, which could allow for populations of *P. armatus* to expand their range northward in years with mild winters.

Tarpon (*Megalops atlanticus*) recruitment to marsh pools and impoundments along the South Carolina coast

Investigators: Garrett Elmo¹, Derek Crane¹, Matthew E. Kimball², and Erin Hosto³

1 - Department of Biology, Coastal Carolina University (SC)

2 - Baruch Marine Field Laboratory, University of South Carolina

3 - Cornell College (IA)

Tarpon (larval and juvenile) will be collected from high marsh pools and impoundments from three estuarine locations along the South Carolina coast: the North Inlet estuary, the Tom Yawkey Wildlife Center, and Kiawah Island. Sampling will consist of pulling a plankton net across one or more fixed transects to sample the surface of the water column, and using a cast net (3 throws) to collect larger nekton at each location. Plankton samples will be put on ice and transported to Baruch Marine Field Laboratory, where larval fishes will be sorted to identify, count, and measure (mm, total length) larval tarpon. Each larval tarpon (i.e., leptocephali) will be categorized based upon developmental stage. Larval density will be calculated by dividing the number of leptocephali collected by the volume of water sampled. Nekton collected with cast nets will be sorted and counted by species in the field, with any larger juvenile tarpon collected will be measured (mm; total, forked, standard length). Water quality (dissolved oxygen, salinity, and temperature) will be collected at each site during each sampling event. Descriptive statistics will be used to examine periods of habitat use at all three sites for both larval and juvenile tarpon during the study period (May through November), and ultimately determine if peak timing of tarpon recruitment and duration of ingress differs among sites.

***Petrolisthes armatus* density and performance along a latitudinal gradient**

Investigator: Jeb Byers

Odum School of Ecology, University of Georgia

Petrolisthes armatus is an invasive tropical crab that has spread throughout the southeastern US in the past decades. Its northern distribution seems to have remained close to North Inlet for many years, perhaps because the crab is limited by low winter temperatures. We have been latitudinally sampling the crab for the past 2 years to establish information on its distribution and abundance, and also to collect crabs for genetic analyses. In particular we wish to examine whether there is any genetic differentiation in the crab within its invasive range. As the northernmost end point in our latitudinal sampling, the North Inlet estuary is an important spot to include in our analyses.

Archaeological investigations of an early to late 18th century house site - testing methods of intrasite patterning detection

Investigators: Heathley Johnson and Karen Smith

South Carolina Institute of Archaeology and Anthropology, University of South Carolina

Archaeological research at site 38GE432 is being conducted to explore the potential for using a metal detector to predict intra-site patterning, as well as to investigate the site itself. A survey of the site using a metal detector on a grid was conducted, with artifact locations mapped with a transit. Artifact distribution maps will then be used to guide the placement of excavation units in order to validate the patterning predicted by the survey. This work will add to the body of knowledge regarding the potential uses of metal detectors as an archaeological survey method, while the resulting archaeological data will expand the knowledge of early to late eighteenth century colonial households in the Georgetown area.

Conservation of diamondback terrapin (*Malaclemys terrapin*) nesting sites in the North Inlet estuary: Using predator vocalization to deter raccoon (*Procyon lotor*) nest predation

Investigators: Colleen Naeger¹ and Scott Parker²

1 - Department of Coastal and Marine Systems Science, Coastal Carolina University (SC)

2 - Department of Biology, Coastal Carolina University (SC)

Diamondback terrapins experience nest predation from foraging raccoons. The purpose of this study is to determine if predator vocalizations deter raccoons from foraging on diamondback terrapin eggs. The hypothesis is that vocalizations from common raccoon predators, such as coyotes will deter raccoon foraging on diamondback terrapin nests. This study will randomly assign predator, non-predator and no vocalization treatments to a series of nest sites. GEN2 GC350 Programmable Call Machines will project vocalizations, while Moultrie® M-880 trail cameras will observe and quantify raccoon foraging behaviors. Expected outcomes are that predator vocalizations will deter raccoon foraging and conserve critical diamondback terrapin nest habitats

Using genetic techniques to understand the mechanism behind the expression of circatidal clocks in an estuarine crab

Investigators: Paola López-Duarte, Taylor Parker, and Leyna Pence

Department of Biological Sciences, University of North Carolina Charlotte

The use of tidal currents to achieve horizontal displacement in the water column is critical to estuarine-ocean migrations for larval and adult forms of countless of species. Previous research has established that swimming activity involved in the selective-tidal stream transport in fiddler crab larvae is under endogenous control. However, the mechanism that allows organisms to keep track and accurately predict the tide, the circatidal clock, is not entirely understood. our goal is to take advantage of recent advances in the identification of “clock genes” to better understand how organisms process information regarding tidal phase and periodicity and how that in turn results in the expression of tidal rhythms (12.4-hour or 24.8-hour cycles).

Physiological plasticity and adaptation to microhabitats within populations of the eastern oyster (*Crassostrea virginica*)

Investigators: Denise Furr, Anna Ivanina, and Adam Reitzel

Department of Biological Sciences, University of North Carolina Charlotte

The eastern oyster is benthic bivalve found on the Atlantic coast, including North Carolina. Environmental variation between tidal zones are ideal for studying ecological factors that affect interspecies variation at local scales. No data are currently available on combined genetics and physiology of oysters from close locations or microhabitats to determine how genetic and environmental factors affect location-specific resilience. Site as well as microhabitat factors are investigated by studying tidal oysters in diverse sites as to energy storage, bioenergetics, immunity expression and genomics. Oysters from different microhabitats should exhibit a different response to the environment, whether these are determined by physiological or genetic adaptation, providing insight into alternative metabolisms and culturing conditions for the shellfish industry.

Characterizing the population genomics of the hemichordate *Saccoglossus kowalevskii*

Investigators: Caitlin Redak, Ken Halanych, and Michael Tassia
Department of Biological Sciences, Auburn University (AL)

We aim to characterize the population genomic structure of acorn worms (*Saccoglossus kowalevskii*) along the US east coast and elucidate any previously unknown relationships between geographically distinct groups. To do this we will collect animals at 6-7 locations and characterize their genomic variation using whole genome sequencing. By utilizing whole genomes we will be able to understand regions of the genome that are under selection and those which are conserved across populations. We also will analyze this dataset for levels of variation that may warrant a redefining of the species, potentially splitting the group into two or more distinct taxa. This project will help inform current and future use of *S. kowalevskii* as a model species.

Developing a behavioral bioindicator of habitat health for coastal salt marshes

Investigator: Blaine D. Griffen
Department of Biology, Brigham Young University (UT)

The goal of this study is to compare herding behavior of sand fiddler crabs across marshes with varying levels of human disturbance to determine whether changes in crab behavior can be predictably used as an indicator of marsh health. At 25-30 marsh sites along the coast between Georgetown and Myrtle Beach that experience varying levels of human disturbance, we will measure herding behavior using a simple experiment following methods of Viscido and Wethey (2002) where a predatory attack is simulated by an individual walking towards a herd. Experiments will be video recorded and later analyzed. In addition, at each site we will assess the level of human disturbance (e.g., distance to roads, infrastructure, amount of foot and road traffic, etc.).

The effects of climate change on *Littoraria irrorata* and *Ilyanassa obsoleta* distribution and movement

Investigators: Samantha McClain and Students Engaged in Aquatic Sciences members
School of the Earth, Ocean, and Environment, University of South Carolina

Two types of marsh snail, the marsh periwinkle (*Littoraria irrorata*) and the eastern mud snail (*Ilyanassa obsoleta*), are common to the North Inlet estuary. Data will be collected concerning their size, distribution, and movement throughout tidal cycles. Using environmental data collected by the NI-WB NERR over several years, the effects of climate change, including water and air temperature, on each species of snail will be revealed and will be used to indicate how climate change is effecting the greater marsh ecosystem. Additionally, this project will provide experience in field research and oceanography for the members of the University of South Carolina undergraduate program Students Engaged in Aquatic Sciences (SEAS).

Analysis of benthic microalgae composition in salt pan and vegetated marsh sediments in the North Inlet estuary, South Carolina

Investigators: Ashley Foulk, Ashley Davis, Abby Davis, and Students Engaged in Aquatic Sciences members
School of the Earth, Ocean, and Environment, University of South Carolina

Benthic microalgae are the foundation of the food chain, but little is known about microalgae communities in the high-salinity, non-vegetated salt pan habitats of the salt marsh. This year, Students Engaged in Aquatic Sciences (SEAS), a program dedicated to providing University of South Carolina (USC) undergraduates the opportunity to conduct research on coastal and oceanic processes, will evaluate the microalgae community in salt pans and adjacent vegetated salt marsh in the North-Inlet estuary. Specifically, USC SEAS members will collect sediment samples from two salt pan and two vegetated marsh locations along the causeway to Clambank Landing, and analyze the 5mm cores using high-performance liquid chromatography. This study will provide insight on how microalgae composition varies between marshes and salt pans, and determine if microalgae communities experience seasonal variation.

Geographic distribution of gelatinous zooplankton in the North Inlet-Winyah Bay estuarine system

Investigators: Joshua Stone¹ and Imani Haley²

1 - Department of Biological Sciences, University of South Carolina

2 - College of Education, University of South Carolina

The North Inlet-Winyah Bay estuarine system is a dynamic system with short retention times and strong currents. In order to determine the distribution of gelatinous zooplankton within the system, we sampled throughout the North Inlet estuary and parts of Winyah Bay over three weeks in June of 2019. We used surface and epi-benthic zooplankton net tows to collect zooplankton samples that will be analyzed and counted over the fall of 2019 for gelatinous zooplankton abundance. The samples were taken across transects from the mouth of the estuary to the heads of the tidal creeks as well as a mid-estuary location across a full tidal cycle. These data will be used to determine the environmental conditions most favorable to gelatinous zooplankton, as well as the total abundance of gelatinous zooplankton throughout the estuarine system

Heart development and molecular characterization of the teleost *Megalops atlanticus*

Investigators: Haley A. Davies¹, Matthew E. Kimball², Aaron J. Adams³, and Frank L. Conlon¹

1 - Integrative Program for Biological & Genome Sciences, University of North Carolina Chapel Hill School of Medicine

2 - Baruch Marine Field Laboratory, University of South Carolina

3 - Bonefish & Tarpon Trust (FL)

The tarpon (*Megalops atlanticus*), a bony, ray-finned fish in the teleost subclass, is a prized game-fish that has been poorly studied. Teleosts comprise the largest subclass of the Actinopterygii class and include 96% of extant fish species. Tarpon fishing for sport is a tourism industry that generates millions of dollars every year for the economies of Central American countries including Costa Rica, Belize, and Panama. In collaboration with the Bonefish and Tarpon Trust and the Belle W. Baruch Institute for Marine and Coastal Sciences at the University of South Carolina, our lab has acquired samples to begin characterization studies on this species. Tarpon are one of very few fish to have an obligatory requirement to breathe atmospheric oxygen for at least some portion of its lifespan. We aim to elucidate the mechanisms underlying the intricacies of the co-development of the heart and swim bladder from an evolutionary perspective by characterizing this non-model organism. As the characterization of non-model organisms becomes more feasible with the decreasing costs of sequencing, supplemental comparisons between these non-model organisms with traditional model systems become more possible. Here, as we begin to study tarpon, we describe the general anatomy with whole-mount skeletal visualization and histology. In addition, we performed the first whole tarpon RNAseq paired with de novo transcriptome assembly as well as preliminary proteomic profiling. Preliminary transcriptome assembly revealed transcripts with sequence homology to known cardiac master regulators, such as *Tbx5*, *Gata4*, and *Nkx2-5*. Future directions will investigate more specific aspects of the cardiac and respiratory functions of tarpon.

The implications of geographic location and intraspecific size variation on *Littoraria irrorata* personality types

Investigators: Christina Salerno and Stephanie Kamel

Department of Biology and Marine Biology, University of North Carolina Wilmington

One of the most abundant and commonly studied invertebrates in southeastern US salt marshes is the marsh periwinkle, *Littoraria irrorata*. Previous research has determined that individual *L. irrorata* exhibit distinct personality types that are consistent over time. This proposed research will further investigate the presence of personality types and examine the effects of snail size, geographic location, refugia presence, and predator presence on individual personality types of *L. irrorata* from four states using a series of experiments. Intraspecific variation of *L. irrorata* personality types can have many ecological implications including predator-prey interactions, community compositions, population stability, and overall marsh health.

Author Index

A

Adams, Aaron J.	34
Allen, Dennis M.	11, 14, 15, 18, 20
Allen, Wendy	13, 21, 27
Atkins, Rebecca	16

B

Baden, John	10, 12, 24
Baker, Julia P.	17
Benitez-Nelson, Claudia	28
Boyer, Emma	21
Brabson, Bill	13, 27
Buck, Tracy	10, 21, 25, 29
Byer, M. D.	24
Byers, Jeb	31
Byrd, Brian	25

C

Capella, Danielle	20
Chestnut, David E.	19
Childress, Michael	29
Clendenin, William	15
Conlon, Frank L.	34
Cothran, Jeremy	22
Currie, Lily	11
Crane, Derek P.	28, 30, 31

D

Daugomah, James	23
Davies, Haley A.	34
Davis, Abby	33
Davis, Ashley	33
Davis, Shannon	20
de Buron, Isaure	18
Denham, Susan	10, 12

E

Ellis, Jean T.	27
Elmo, Garrett	28, 30, 31

F

Fangman, Isabel	
Folk, Brooks	24

F (cont.)

Foulk, Ashley	33
Fournier, Hayley	15
Funk, Thomas	22
Furr, Denise	32

G

Gerard, Patrick	29
Gorr, Alex	20
Griffen, Blaine D.	12, 33

H

Halanych, Ken	33
Haley, Imani	34
Hawk, Jennifer	25
Hawkins, Daniel	16
Harding, Juliana M.	14, 16, 18, 19, 20, 25
Hosto, Erin	31
Houser, Kyle	20

I

Ide, Melissa	22, 24
Ivanina, Ann	32

J

Jachowski, David	29
Jefferson, William H.	13, 24
Jodice, Patrick G.R.	27
Johnson, Heathley	32
Johnson, Sharleen	19
Jung, A.	24

K

Kamel, Stephanie	34
Keim, Richard	18
Kenny, Paul D.	11, 21
Kerr, Cindy	13
Kessee, Jennifer	24
Key, Peter	23
Kibler, Krystyn	26
Kimball, Matthew E.	11, 12, 15, 20, 26, 28, 30, 31, 34
Kingston, Alexandra	23
Knotts, Eilea	22

K (cont.)

Knowles, Amber	24
Konsek, Marlene	13, 27
Krask, Julie L.	10, 12, 21, 25, 29
Kyle, Dennis	18

L

LaPlaca, Stephanie	14
Lawson, Abigail J.	27
Lee, Timothy	16
Little, Pete	13
Little, Sandy	13
Paola López-Duarte, Paola	32
Lucia, Rebecca	23

M

Markezich Allan L.	12
McCabe, Kelly	28
McClain, Samantha	33
Miller, Gwen	24
Morris, James T.	17, 18, 23, 24

N

Naeger, Colleen	32
-----------------------	----

O

O'Brien, Danielle	17
O'Halloran, Thomas L.	13
Osenberg, Craig	16

P

Parker, Scott	20, 32
Parker, Taylor	32
Pence, Leyna	32
Pinckney, James L.	22, 26, 33
Pfirmsmann, Bruce	11, 12, 15
Plunket, Jennifer	13, 17, 21, 27
Popp, Teresa	31
Porter, Dwayne E.	22, 24

R

Rachlin, Joseph	11
Rainwater, Thomas R.	27
Ramage, Dan	24
Redak, Caitlin	333
Reitzel, Adam	16, 32

S

Salerno, Christina	34
Sanger, Denise	19
Sheremet, Vitalii A.	19
Shutt, Lee	24
Smith, Erik M.	10, 12, 13, 19, 20, 21, 25, 26, 28, 29
Smith, Karen	32
Snyder, Maeve	14
Speiser, Daniel I.	23
Stalter, Richard	10, 11, 24
Starosta, A.	24
Stevens, Baker	10, 12, 21, 25, 29
Stone, Joshua	15, 34
Strosnider, William	11
Sundberg, Karen	17, 18, 23

T

Tasker, Travis	11
Tassia, Michael	33
Thomas, Beth	15
Torres, Raymond	18
Tweel, Andrew	19

V

Van den Hurk, Peter	14
Voulgaris, George	13

W

Waits, Lisette	29
Walters, Keith	22
Wanderley, Barbara	17
Wang, Cuizhen (Susan)	27, 29
Weinstein, John	14
West, Blaine	23
White, Scott	20
Wiggers, Ernie	29
Wilber, Dara	31
Wilker, Jonathan	21
Wilson, Alicia	15
Wilson, Amanda L.	29
Wirth, Ed F.	19

Y

Young, Rob	10
------------------	----

Z

Zaruba, Kendra	111
----------------------	-----