



INDUSTRIAL
MATHEMATICS
INSTITUTE

2003:21

Order of convergence of minmod-
type schemes

B. Popov and O. Trifonov

IMI

Preprint Series

Department of Mathematics
University of South Carolina

Order of Convergence of Minmod-Type Schemes

Bojan Popov* and Ognian Trifonov†

December 12, 2003

Abstract

A class of non-oscillatory numerical methods for solving nonlinear scalar conservation laws in one space dimension is considered. Non-oscillatory schemes are based on minmod limiters and the standard second order representatives are the staggered Nessyahu-Tadmor scheme and the usual TVD2 scheme. It is well known that the L_p -error of monotone finite difference methods for the linear advection equation is exactly $1/2$ for initial data in $W^1(L_p)$, $1 \leq p \leq \infty$. For a second or higher order non-oscillatory schemes very little is known because they are nonlinear even for the simple advection equation. In this paper, in the case of a linear advection equation with monotone initial data, it is shown that the order of the L_2 -error for the standard second order minmod-type schemes is at least $5/8$ in contrast to the exact $1/2$ order for any formal first order scheme.

1 Introduction

We are interested in the scalar hyperbolic conservation law

$$(1) \quad \begin{cases} u_t + f(u)_x = 0, & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = u^0(x), & x \in \mathbb{R}, \end{cases}$$

where f is a given flux function. In recent years, there has been enormous activity in the development of the mathematical theory and in the construction of numerical methods for (1). Even though the existence-uniqueness theory of weak solutions is complete, there are many numerically efficient methods for which the questions of convergence and error estimates are still open. For example, there are many non-oscillatory schemes based on the minmod limiters which are numerically robust, at least in many numerical tests, but theoretical results about convergence and error estimates are still missing [1, 4, 5, 15].

In this paper, we consider a class of the so-called Godunov-type schemes for solving (1). There are two main steps in such schemes: evolution and projection. In the

*Department of Mathematics, Texas A&M University, College Station, TX 77845, USA, popov@math.tamu.edu.

†Department of Mathematics, University of South Carolina, Columbia, SC 29208, USA, trifonov@math.sc.edu. Supported by the NSF DMS Grant No. 9970455.

original Godunov scheme, the projection is onto piecewise constant functions – the cell averages. In the general Godunov-type method, the projection is onto piecewise polynomials. To determine the properties of these schemes it is necessary to study the properties of the projection operator. We limit our attention to the case of piecewise linear projection based on cell averages using minmod limiters for the slope reconstruction and we call such a scheme *minmod-type*. For example, the Nessyahu-Tadmor(NT) scheme [13] is of minmod-type and it is based on staggered evolution, other examples include the second order non-oscillatory central schemes with non-staggered grids given in [6, 7], and the UNO and TVD2 schemes in [4]. Theoretical results about convergence of such schemes to the entropy solution, or error estimates, are still missing. In most cases, the authors give a variation bound for such a scheme which is enough to conclude that the method converges to a weak solution, see [9]. In the case of linear flux, the conservation law (1) is the usual linear advection equation. The theory of *linear* numerical schemes for the linear advection equation was developed in [2]. It is shown that if a linear finite difference scheme is of formal order μ and L_2 -stable, then the order of convergence in L_2 is exactly $\frac{s\mu}{\mu+1}$ for initial data in $W^s(L_2)$. The case $1 \leq p \leq \infty$, $p \neq 2$ was also considered but the theory is more complicated, see [2] for details. Hence, the order of convergence increases with the formal order of the linear scheme. In the case of $BV(\mathbb{R})$ initial data, Tang and Teng show in [14] that all monotone schemes (a special case of first order schemes used in conservation laws) are exactly 1/2 order accurate in L_1 . Therefore, these schemes cannot be very accurate in computations. From numerical point of view, the formal high order non-oscillatory schemes developed in [4, 5, 13, 7, 12] are much better than any monotone scheme but theoretically there was no rigorous result confirming the numerical evidence even in the case of a linear advection equation. The reason is that most high order schemes used in conservation laws are based on minmod limiters and are nonlinear even in the simplest case of linear advection. Therefore, the results in [2] are not valid for any non-oscillatory scheme based on limiters. The only known result was that a non-oscillatory total variation bounded scheme is at least 1/2 order accurate in L_1 . Recently, we showed in [10] that the standard second order schemes (NT or TVD2) are at least 1/2 order accurate in L_2 for any initial data in $W^1(L_2)$. In both cases, the order of convergence proven for the second order schemes is the same as the order for the first order (for example monotone) schemes. In general, it was not clear how to prove better error estimates for second or higher order non-oscillatory or simply nonlinear schemes even in the case of a scalar linear advection equation. The difficulty in proving better estimates is that even though the minmod-type schemes are formally second order they are known to preserve only first order smoothness of the initial data: (i) the total variation for initial data in $BV(\mathbb{R})$; (ii) a discrete l_2 of the first derivative for initial data in $W^1(L_2(\mathbb{R}))$ (see [10]). Hence, all existing error estimates were proven in the same way as for monotone schemes because they also preserve the above functionals. Here, we develop a new approach for proving better error estimates for second order schemes in the case of the linear advection equation with initial data in $W^1(L_2)$. Namely, we measure how well a scheme approximates a discrete (l_2) norm of the first derivative in time. In our earlier paper [10], we showed that some first (Godunov and LxF) and second order (NT and TVD2) schemes diminish the the l_2 norm of the first

derivative in time and based on that we derived an error estimate with a rate $1/2$ for any initial data in $W^1(L_2)$. Here, we prove that this decay for a second order scheme is different (smaller) than the decay of any first order scheme in the case of a monotone initial data. The new decay estimate allows us show that the error of the standard non-oscillatory second order scheme is at least $5/8$ for any monotone initial data in $W^1(L_2)$. The restriction to monotone initial data is natural because the minmod-type schemes reduce to a formal first order scheme at any local extremum. The paper is organized as follows. In section 2, we describe the standard second order minmod-type schemes (NT and TVD2) in the case of a linear advection equation and give a new decay estimate (see Lemma 1) for a discrete norm of the first derivative of the numerical solution in time. In Section 3, we present our main result: minmod-type schemes have better convergence rate than first order schemes for monotone initial data. In the appendix, we give the proof of Lemma 1.

2 Stability of non-oscillatory minmod-type schemes

In this section, we are concerned with non-oscillatory differencing approximations to the linear advection equation

$$(2) \quad \begin{cases} u_t + au_x = 0, & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = u^0(x), & x \in \mathbb{R}, \end{cases}$$

The prototypes of all high-order non-oscillatory schemes are the second order schemes based on a piecewise linear spacial reconstruction. We restrict our attention to the standard representatives: the usual TVD2 scheme, see for example [4], and the staggered Nessyahu-Tadmor (NT) scheme [13]. In the case of a linear flux, $f(u) = au$, and uniform space and time meshes, both schemes reduce to the same iterative relation between the sequences of new and old cell averages (note that the NT scheme alternates between two uniform grids). For simplicity, we only consider the case $a \geq 0$ with the other case being analogous. Then, the relationship between *new* and *old* cell averages is

$$(3) \quad w'_j = \alpha w_{j-1} + (1 - \alpha)w_j + \frac{1}{2}\alpha(1 - \alpha) (m(\delta_{j-1}, \delta_j) - m(\delta_j, \delta_{j+1})),$$

where $m(a, b)$ stands for the minmod limiter

$$(4) \quad m(a, b) \equiv \text{MinMod}(a, b) := \frac{1}{2}(\text{sgn}(a) + \text{sgn}(b)) \cdot \min(|a|, |b|),$$

and $\delta_j := w_j - w_{j-1}$, $j \in \mathbb{Z}$. Here we use $\{w_j\}$ for the averages of the numerical solution v at the old time step $t_n = n\Delta t$ and $\{w'_j\}$ for the averages of the numerical solution v at the new time step $t_{n+1} = (n+1)\Delta t$. Let $x_j := j\Delta x$, $j \in \mathbb{Z}$, and $\lambda := \frac{\Delta t}{\Delta x}$. In the case of the NT scheme (see [10]), we have that

$$w_j := \frac{1}{\Delta x} \int_{x_{j-1/2}}^{x_{j+1/2}} v(x, t_n) dx, \quad w'_{j+1} := \frac{1}{\Delta x} \int_{x_j}^{x_{j+1}} v(x, t_{n+1}) dx, \quad \alpha := \frac{1}{2} + \lambda a,$$

and the CFL condition reduces to $\lambda a \leq \frac{1}{2}$. In the case of the TVD2 scheme (see section 7 of [4]), we have that

$$w_j := \frac{1}{\Delta x} \int_{x_{j-1/2}}^{x_{j+1/2}} v(x, t_n) dx, \quad w'_j := \frac{1}{\Delta x} \int_{x_{j-1/2}}^{x_{j+1/2}} v(x, t_{n+1}) dx, \quad \alpha := \lambda a,$$

and the CFL condition reduces to $\lambda a \leq 1$. Note that both CFL conditions imply the same restriction on α

$$(5) \quad 0 \leq \alpha \leq 1.$$

and the sequences of cell averages of the numerical solution generated by the NT scheme for one value of a are the same as the sequences of the TVD2 scheme with a different a but the same initial data if they both result in the same value for α . Hence, the numerical stability of each scheme is the same over the class of all linear flux functions $f(u) = au$.

Let us denote the numerical solution of either scheme (NT or TVD2) at time step t_n with v^n , $v^n := v(\cdot, t_n)$, and its cell averages with v_j^n . Note that the cell averages of v^0 are equal to the cell averages of the initial condition u^0 : $v_j^0 := u_j^0$, $j \in \mathbb{Z}$, and the numerical solution $v(\cdot, t_n)$ is a linear function on each cell $I_j := (x_{j-1/2}, x_{j+1/2})$

$$(6) \quad v^n|_{I_j} = v_j^n + (x - x_j) \frac{1}{\Delta x} m(v_{j+1}^n - v_j^n, v_j^n - v_{j-1}^n).$$

Note that in the NT case the above formula is valid only for even n because for odd n we have a similar piecewise linear reconstruction but on the staggered grid, i.e., we have a minmod-type reconstruction on $I_j := (x_j, x_{j+1})$, $j \in \mathbb{Z}$.

One can define a global numerical solution v corresponding to the initial data u^0 in the following way. Let $v(x, t)$ be a right-continuous function in t such that, for each $n = 0, \dots, N-1$, v is the solution of

$$(7) \quad \begin{cases} u_t^n + au_x^n = 0, & (x, t) \in \mathbb{R} \times (t_n, t_{n+1}), \\ u(x, 0) = v^n(x), & x \in \mathbb{R}, \end{cases}$$

Note that v is uniquely determined by the functions $\{v^n\}_{n=0}^{N-1}$, where v^n is the minmod piecewise linear reconstruction defined in (6) from the cell averages of $v(\cdot, t_n^-)$ with $v(\cdot, 0^-) := u^0$. In the TVD2 case, we have the same uniform grid for each step, and in the NT case we alternate between two staggered uniform grids. In both cases, we end up with the same type of iterative relation between the cell averages of the numerical solution, see (3), just the value of α is different.

We now present a new stability property for a numerical solution which satisfies (3). The new averages $v_j^{n+1} := w'_j$, can be written in terms of the old averages $v_j^n := w_j$ and the old jumps $\delta_j^n := v_j^n - v_{j-1}^n$, $j \in \mathbb{Z}$, in the following way

$$(8) \quad v_j^{n+1} = \alpha v_{j-1}^n + (1 - \alpha) v_j^n + \frac{1}{2} \alpha (1 - \alpha) (m(\delta_{j-1}^n, \delta_j^n) - m(\delta_j^n, \delta_{j+1}^n))$$

for $n \geq 0$. Using that, we derive the formula for the sequence of new jumps $\{\delta_j^{n+1}\}$ in terms of the old ones $\{\delta_j^n\}$

$$(9) \quad \delta_j^{n+1} = \alpha \delta_{j-1}^n + (1 - \alpha) \delta_j^n - \beta m(\delta_{j-2}^n, \delta_{j-1}^n) + 2\beta m(\delta_{j-1}^n, \delta_j^n) - \beta m(\delta_j^n, \delta_{j+1}^n),$$

where $\beta := \frac{1}{2}\alpha(1 - \alpha)$.

In our previous paper [10], we proved that the the iterative formula (9) does not increase the l_2 norm of the jumps of the numerical solution in time. That is,

$$(10) \quad \sum_j (\delta_j^{n+1})^2 - \sum_j (\delta_j^n)^2 \leq 0$$

for all $n = 0, 1, \dots$. The above stability result can be easily verified for any monotone scheme because in that case the new cell averages are a convex combination of the old ones. For a first order scheme (including all monotone schemes) the difference $\sum_j (\delta_j^n)^2 - \sum_j (\delta_j^{n+1})^2$ is of order $\sum_j (\delta_j^n - \delta_{j-1}^n)^2$. This reflects the diffusive nature of the first order approximation. For the exact solution, there is no decay in time of any shift invariant norm but for a first order numerical solution we expect, and it is easy to show that for the Godunov and LxF schemes, the following numerical viscosity in the decay of the first derivative l_2 -norm

$$\sum_j (\delta_j^n)^2 - \sum_j (\delta_j^{n+1})^2 = O\left(\sum_j (\delta_j^n - \delta_{j-1}^n)^2\right) = O\left(\sum_j (v_{j+1}^n - 2v_j^n + v_{j-1}^n)^2\right).$$

Here, we derive an improved stability estimate for the second order minmod-type schemes given in (8). Namely, we give an exact estimate for the time decay of the discrete l_2 -norm in the case of a monotone initial condition. We claim

Lemma 1. *Let u^0 be any monotone function and $\{v^n\}$, $n = 0, 1, \dots$, be the sequence approximate solutions v^n defined by (8). Then, we have*

$$(11) \quad C \sum_j (\Delta^2 \delta_j^n)^2 \geq \sum_j (\delta_j^n)^2 - \sum_j (\delta_j^{n+1})^2 \geq c \sum_j (\Delta^2 \delta_j^n)^2,$$

with $C = 3\beta$ and $c = \beta^3/4$, where $\beta = \frac{1}{2}\alpha(1 - \alpha)$.

The proof of this lemma is given in the appendix. We want to point out the main difference: for a first order scheme, we have a decay for $\sum_j (\delta_j^n)^2 - \sum_j (\delta_j^{n+1})^2$ of order $\sum_j (v_{j+1}^n - 2v_j^n + v_{j-1}^n)^2$, and for a second order scheme we get $\sum_j (v_{j+1}^n - 3v_j^n + 3v_{j-1}^n - v_{j-2}^n)^2$ by Lemma 1. We will use this decay estimate in the next section to derive our main result.

3 Error estimates for second order schemes

In this section we present our main result. Namely, we will show that minmod-type schemes have a better convergence rate than first order schemes for any monotone initial condition $u^0 \in W^1(L_2)$. Let us first recall some results for first order linear schemes (see Theorem 4.2 and Theorem 4.4 in [2]).

Theorem 2. *Let $u^0 \in W^1(L_2)$, u be the exact solution of (2), and v be a numerical solution consistent with (2) and exactly first order accurate. If the space and time meshes*

are $h := \Delta x$ and Δt with $\Delta t/h = \lambda = \text{const}$, then there exists a constant $C > 0$ which depends on the final time $T > 0$ such that

$$\|u(\cdot, T) - v(\cdot, T)\|_{L_2} \leq Ch^{1/2} \|u^0\|_{W^1(L_2)}.$$

Moreover, the above estimate is optimal over the class $W^1(L_2)$. That is, there exists a constant $c > 0$ such that

$$(12) \quad \sup\{\|u(\cdot, T) - v(\cdot, T)\|_{L_2} : u^0 \in \hat{C}^\infty, \|u^0\|_{W^1(L_2)} \leq 1\} \geq ch^{1/2},$$

where \hat{C}^∞ is the set of all functions with Fourier transform in $C_0^\infty(\mathbb{R})$.

The above result is also valid for any monotone scheme except in the trivial case of pure translation which corresponds to $\alpha = 0$ or $\alpha = 1$ here (see Remark 1 in [14] and Section 3.3 in [2]) because the numerical solution in these two cases is the same as the exact solution. Using the linearity of the problem and the numerical scheme, it is easy to show

Corollary 3. *The convergence rate for any linear exactly first order scheme is not better than 1/2 for the class of all monotone initial conditions $u^0 \in W_{loc}^1(L_2(\mathbb{R}))$ with $\frac{du^0}{dx} \in L_2(\mathbb{R})$. That is, an estimate of the type*

$$\|u(\cdot, T) - v(\cdot, T)\|_{L_2} \leq Ch^\gamma |u^0|_{W^1(L_2)}$$

for all monotone u^0 is not possible for $\gamma > 1/2$.

The seminorm $|u^0|_{W^1(L_2)}$ is a natural replacement of the full norm in estimates for equations with monotone initial conditions. It is also important to note that all results above, including the lower bound (12), hold for initial conditions u^0 which have first derivatives with compact support. That is, the error estimate and the lower bound in Theorem 2 do not depend on the infinite space domain. In the case of the Godunov or LxF schemes one can give an example of a monotone function with a derivative with compact support (similar to the construction in [14]) which realizes the lower bound (12). Hence, there are no ‘‘log’’ effects and the rate of convergence is exactly 1/2 for these two schemes.

The following theorem is our main result. That is, the standard second order minmod-type schemes, NT and TVD2, have a better rate of convergence than the rate of any formal first order (including monotone) scheme.

Theorem 4. *Let $u^0 \in W^1(L_2)$ be monotone, u be the exact solution of (2), and v be the numerical solution generated by a standard minmod-type scheme (NT or TVD2), see (6) and (8). If the CFL condition $0 < \alpha < 1$ is satisfied, $h := \Delta x$, $t_n = n\Delta t$, $0 \leq n \leq N$, and $T = N\Delta t$, we have*

$$(13) \quad \|u(T, \cdot) - v(T, \cdot)\|_{L^2} \leq Ch^{5/8} |u^0|_{W^1(L_2)},$$

where C is a constant which depends only on the final time T and α .

Proof. The error estimate is based on a refinement of the dual argument in [10] and the new stability result Lemma 1. In the proof, C will be an absolute constant that can be different at different places. It is enough to prove the estimate for initial condition u^0 which has a derivative with compact support. The general result follows by standard arguments using the local dependence of the exact and the numerical solutions.

Let $e(x, t) := u(x, t) - v(x, t)$ be the difference between the global numerical solution v defined in (7) and the exact solution u , and $E(x, t) := \int_{-\infty}^x e(s, t) ds$. Note that $u^0 - v^0$ is zero for $x \in (-\infty, M)$, for some $M \in \mathbb{R}$. Therefore E is well defined for all $(x, t) \in \mathbb{R} \times (0, T)$. We have that E also satisfies (2) for $n\Delta t \leq t < (n+1)\Delta t$ with initial data $\int_{-\infty}^x u(s, n\Delta t) - v^n(s) ds$, $n = 0, 1, \dots, N-1$. For a function $g \in L^1(\mathbb{R})$, we define a *minus one* norm in the following way

$$(14) \quad \|g\|_{-1,2} := \left\| \int_{-\infty}^{\cdot} g(s) ds \right\|_{L^2(\mathbb{R})}.$$

It is easy to verify that for any $\tau \in \mathbb{R}$

$$(15) \quad \|S_\tau g\|_{-1,2} = \|g\|_{-1,2},$$

where S_τ is the shift operator $S_\tau g(\cdot) := g(\cdot - \tau)$.

Let us denote the piecewise linear minmod reconstruction (6) with P_h . That gives $v^n = P_h v(\cdot, t_n -)$ for $n = 0, 1, \dots, N$, and $u(\cdot, t) = S_t(u^0)$ for any $t \geq 0$. Recall that $T = N\Delta t$. Then, we have the representations $u(T, \cdot) = (S_{a\Delta t})^N u^0$ and $v(T, \cdot) = v^N = P_h(S_{a\Delta t} P_h)^N u^0$. We have

$$\|e(T, \cdot)\|_{-1,2} = \|(S_{a\Delta t})^N u^0 - P_h(S_{a\Delta t} P_h)^N u^0\|_{-1,2},$$

and by the triangle inequality we obtain

$$(16) \quad \begin{aligned} \|e(T, \cdot)\|_{-1,2} &\leq \|(S_{a\Delta t})^N u^0 - (S_{a\Delta t} P_h)^N u^0\|_{-1,2} \\ &\quad + \|P_h(S_{a\Delta t})^N u^0 - (S_{a\Delta t} P_h)^N u^0\|_{-1,2}. \end{aligned}$$

Using (15) in (16), we obtain

$$(17) \quad \|e(T, \cdot)\|_{-1,2} \leq \|(S_{a\Delta t})^{N-1} u^0 - P_h(S_{a\Delta t} P_h)^{N-1} u^0\|_{-1,2} + \|v^N - v(\cdot, t_N -)\|_{-1,2}.$$

Let e^n be the difference between the exact and the numerical solution at time $t_n = n\Delta t$. That is, $e_n := ((S_{a\Delta t})^n - P_h(S_{a\Delta t} P_h)^n) u^0$, $n = 0, 1, \dots, N$. Then (17) is equivalent to

$$(18) \quad \|e^N\|_{-1,2} \leq \|e^{N-1}\|_{-1,2} + \|P_h v(\cdot, t_N -) - v(\cdot, t_N -)\|_{-1,2},$$

and applying (18) for $n = N, N-1, \dots, 1$, we get

$$(19) \quad \|e^N\|_{-1,2} \leq \sum_{n=0}^N \|P_h v(\cdot, t_n -) - v(\cdot, t_n -)\|_{-1,2}$$

We now estimate the minus one norm of $P_h v(\cdot, t_n -) - v(\cdot, t_n -)$ in terms of the usual L_2 norm. In order to simplify the notation, we will use $v^{n,-} := v(\cdot, t_n -)$. We have

$$\begin{aligned}
\|v^n - v^{n,-}\|_{-1,2}^2 &= \int_{\mathbb{R}} \left(\int_{-\infty}^x (v^n(s) - v^{n,-}(s)) ds \right)^2 dx \\
(20) \qquad \qquad \qquad &= \sum_{j \in \mathbb{Z}} \int_{x_{j-1/2}}^{x_{j+1/2}} \left(\int_{x_{j-1/2}}^x (v^n - v^{n,-})(s) ds \right)^2 dx,
\end{aligned}$$

where we use in (20) that $v^n = P_h v^{n,-}$ is a conservative approximation of v^n , i.e., $\int_{x_{j-1/2}}^{x_{j+1/2}} (v^n - v^{n,-})(s) ds = 0$ for all $j \in \mathbb{Z}$. Note that in the case of the NT scheme we need to use conservation over the staggered intervals (x_j, x_{j+1}) , $j \in \mathbb{Z}$, for odd n . After applying the Cauchy-Schwarz inequality in (20), we obtain

$$\begin{aligned}
\|v^n - v^{n,-}\|_{-1,2}^2 &\leq \sum_{j \in \mathbb{Z}} \int_{x_{j-1/2}}^{x_{j+1/2}} (x - x_{j-1/2}) \int_{x_{j-1/2}}^x (v^n(s) - v^{n,-}(s))^2 ds dx \\
&\leq \sum_{j \in \mathbb{Z}} \int_{x_{j-1/2}}^{x_{j+1/2}} (x - x_{j-1/2}) dx \int_{x_{j-1/2}}^{x_{j+1/2}} (v^n(s) - v^{n,-}(s))^2 ds \\
&= \frac{h^2}{2} \|v^n - v^{n,-}\|_{L_2(\mathbb{R})}^2.
\end{aligned}$$

Therefore, the error estimate (19) reduces to

$$(21) \qquad \|e^N\|_{-1,2} \leq \frac{h}{\sqrt{2}} \sum_{n=0}^N \|v^n - v^{n,-}\|_{L_2(\mathbb{R})}.$$

We now estimate the term $\|v^n - v^{n,-}\|_{L_2(\mathbb{R})}$. Because the minmod operator P_h is non-linear, we will use an intermediate approximation \tilde{P}_h defined in the following way

$$\tilde{P}_h v^{n,-}|_{I_j} = v_j^n + \frac{x - x_j}{h} (v_{j+1}^n - v_j^n),$$

for $j \in \mathbb{Z}$. It is a straightforward computation that

$$(22) \qquad \|P_h v^{n,-} - \tilde{P}_h v^{n,-}\|_{L_2(\mathbb{R})} \leq Ch^{1/2} \|\Delta^2 v_j^n\|_{l_2}.$$

Hence, we have the estimate

$$(23) \qquad \|v^n - v^{n,-}\|_{L_2(\mathbb{R})} \leq Ch^{1/2} \|\{\Delta^2 v_j^n\}\|_{l_2} + \|\tilde{P}_h v^{n,-} - v^{n,-}\|_{L_2(\mathbb{R})}.$$

\tilde{P}_h is a linear operator based on local averages. The local approximation properties of \tilde{P}_h were analyzed in [8]. Using that result, it is easy to show that

$$(24) \qquad \|\tilde{P}_h g - g\|_{L_2(\mathbb{R})} \leq Ch^2 \|g''\|_{L_2(\mathbb{R})}, \quad \text{for } g \in W^2(L_2(\mathbb{R})),$$

$$(25) \qquad \|\tilde{P}_h g\|_{L_2(\mathbb{R})} \leq C \|g\|_{L_2(\mathbb{R})} \quad \text{for } g \in L_2(\mathbb{R}).$$

Then, by standard approximation theory arguments (see Chapter 7 in [3]), we derive

$$\|\tilde{P}_h g - g\|_{L_2(\mathbb{R})} \leq C\omega_2(g, h)_{L_2(\mathbb{R})}$$

for any $g \in L_2(\mathbb{R})$. Here $\omega_2(g, h)_{L_2(\mathbb{R})}$ is the second modulus of smoothness of g in L_2 . Using the above estimate with $g := v^{n,-}$, we obtain

$$(26) \quad \|\tilde{P}_h v^{n,-} - v^{n,-}\|_{L_2(\mathbb{R})} \leq C\omega_2(v^{n,-}, h)_{L_2(\mathbb{R})} = C\omega_2(v^{n-1}, h)_{L_2(\mathbb{R})}$$

because $v^{n,-} = S_{\alpha\Delta t} v^{n-1}$ and the modulus of smoothness $\omega_2(v^{n-1}, h)_{L_2(\mathbb{R})}$ is shift invariant. In general, it is not easy to compute $\omega_2(g, h)_{L_2(\mathbb{R})}$ for a general g but in the case of the piecewise linear function v^n it is not hard to show that for any $n = 0, 1, \dots, N$, we have

$$(27) \quad C_1 h^{1/2} \|\{\Delta^2 v_j^n\}\|_{l_2} \leq \omega_2(v^n, h)_{L_2(\mathbb{R})} \leq C_2 h^{1/2} \|\{\Delta^2 v_j^n\}\|_{l_2}.$$

Finally, we use (27) and (26) in (23) and obtain

$$(28) \quad \|v^n - v^{n,-}\|_{L_2(\mathbb{R})} \leq Ch^{1/2} (\|\{\Delta^2 v_j^{n-1}\}\|_{l_2} + \|\{\Delta^2 v_j^n\}\|_{l_2}).$$

We now go back and use (28) in the error estimate (19) to derive

$$(29) \quad \|e^N\|_{-1,2} \leq Ch^{3/2} \sum_{n=0}^N \|\{\Delta^2 v_j^n\}\|_{l_2} = Ch^{3/2} \sum_{n=0}^N \|\{\Delta \delta_j^n\}\|_{l_2},$$

here we recall that $\delta_j^n = v_j^n - v_{j-1}^n$. Up to this point, our arguments are very similar to the ones in our previous paper [10]. The only difference is that here, in the estimate for the negative norm (29), we use $\|\{\Delta \delta_j^n\}\|_{l_2}$ instead of $\|\{\delta_j^n\}\|_{l_2}$ that was used in [10]. A new idea here will be the use of the improved l_2 stability, Lemma 1, to derive a better error estimate than before. Let us recall the lower bound in the statement of Lemma 1. For any $n = 0, \dots, N-1$, we have

$$(30) \quad \sum_j (\delta_j^n)^2 - \sum_j (\delta_j^{n+1})^2 \geq c \sum_j (\Delta^2 \delta_j^n)^2,$$

with $c = \beta^3/4$, where $\beta = \frac{1}{2}\alpha(1-\alpha)$. We sum (30) for $n = 0, \dots, N-1$ and obtain

$$(31) \quad c \sum_{n=0}^N \sum_j (\Delta^2 \delta_j^n)^2 \leq \sum_j (\delta_j^0)^2 - \sum_j (\delta_j^N)^2.$$

Recall that the CFL condition is $0 < \alpha < 1$. In the trivial case $\alpha(1-\alpha) = 0$, not covered by our theorem, we have a pure translation and the total error is equal to the error of the first step, i.e., the order of the error is $O(h)$. So, without loss of generality, we can assume that $0 < \alpha < 1$. Then, we have that $c \neq 0$ and we derive from (31) the following

$$(32) \quad \sum_{n=0}^N \sum_j (\Delta^2 \delta_j^n)^2 \leq C \sum_j (\delta_j^0)^2,$$

where C is an absolute constant for any fixed $0 < \alpha < 1$ ($C = 1/c = 4\beta^{-3}$). Using (32), we will now estimate $\sum_{n=0}^N \|\{\Delta\delta_j^n\}\|_{l_2}$. Because the support of $\frac{du^0}{dx}$ is finite and the numerical solution has a finite domain of dependence, we have that the set $\Lambda_n := \{j : \delta_j^n \neq 0\}$ is finite for all $n = 0, 1, \dots, N$. Using Abel's transform and the finite support of $\{\delta_j^n\}$, we obtain

$$\sum_j (\Delta\delta_j^n)^2 = - \sum_j \delta_j^n (\delta_{j+1}^n - 2\delta_j^n + \delta_{j-1}^n).$$

Hence, we have

$$\sum_j (\Delta\delta_j^n)^2 \leq \sum_j |\delta_j^n| |\delta_{j+1}^n - 2\delta_j^n + \delta_{j-1}^n| \leq \|\{\delta_j^n\}\|_{l_2} \|\{\Delta^2\delta_j^n\}\|_{l_2}.$$

Using that $\|\{\delta_j^n\}\|_{l_2} \leq \|\{\delta_j^0\}\|_{l_2}$, we derive

$$(33) \quad \|\{\Delta\delta_j^n\}\|_{l_2} \leq \|\{\delta_j^0\}\|_{l_2}^{1/2} \|\{\Delta^2\delta_j^n\}\|_{l_2}^{1/2}.$$

Therefore, we have the estimates

$$(34) \quad \begin{aligned} \sum_{n=0}^N \|\{\Delta\delta_j^n\}\|_{l_2} &\leq \|\{\delta_j^0\}\|_{l_2}^{1/2} \sum_{n=0}^N \|\{\Delta^2\delta_j^n\}\|_{l_2}^{1/2} \\ &\leq \|\{\delta_j^0\}\|_{l_2}^{1/2} (N+1)^{3/4} \left(\sum_{n=0}^N \|\{\Delta^2\delta_j^n\}\|_{l_2}^2 \right)^{1/4}, \end{aligned}$$

where we use Hölder's inequality to derive (34). We use our stability estimate (32) in (34) and conclude

$$(35) \quad \sum_{n=0}^N \|\{\Delta\delta_j^n\}\|_{l_2} \leq (N+1)^{3/4} \|\{\delta_j^0\}\|_{l_2}.$$

From Lemma 4 in [10], we have that $\|\{\delta_j^0\}\|_{l_2} \leq h^{1/2} |u^0|_{W^1(L_2(\mathbb{R}))}$. Hence,

$$(36) \quad \sum_{n=0}^N \|\{\Delta\delta_j^n\}\|_{l_2} \leq CN^{3/4} h^{1/2} |u^0|_{W^1(L_2(\mathbb{R}))}.$$

We use (36) in (29) and derive the final minus one norm error estimate

$$(37) \quad \|e^N\|_{-1,2} \leq C(Nh)^{3/4} h^{5/4} |u^0|_{W^1(L_2(\mathbb{R}))} \leq Ch^{5/4} |u^0|_{W^1(L_2(\mathbb{R}))},$$

where in the last inequality C depends on T and α . Similar to [10], we will now interpolate between the negative norm estimate (37) and the discrete W_2^1 -stability of the numerical scheme (10). Recall that $e^N = u(\cdot, T) - v^N$ is not in $W^1(L_2(\mathbb{R}))$. Hence, we approximate v^N with $\tilde{v}^N \in W^1(L_2(\mathbb{R}))$, see the last displayed equation before (54) in [10]. The function \tilde{v}^N has the properties (see (54), (55), and (56) in [10])

$$(38) \quad \|\tilde{v} - v^N\|_{-1,2} \leq Ch^2 |u^0|_{W^1(L_2(\mathbb{R}))},$$

$$(39) \quad \|\tilde{v} - v^N\|_{L^2(\mathbb{R})} \leq Ch|u^0|_{W^1(L_2(\mathbb{R}))},$$

and

$$(40) \quad \|\tilde{v}\|_{W^1(L_2(\mathbb{R}))} \leq |u^0|_{W^1(L_2(\mathbb{R}))}.$$

Let $\tilde{e} := u(T, \cdot) - \tilde{v}$. Then

$$\|\tilde{e}\|_{-1,2} \leq \|e^N\|_{-1,2} + \|\tilde{v} - v^N\|_{-1,2}.$$

Using (37) and (38), we obtain

$$(41) \quad \|\tilde{e}\|_{-1,2} \leq Ch^{5/4}|u^0|_{W^1(L_2(\mathbb{R}))}.$$

Let $\tilde{E}(x) = \int_{-\infty}^x \tilde{e}(s) ds$. Then we have $\|\tilde{E}\|_{L_2} = \|\tilde{e}\|_{-1,2}$, $\|\tilde{E}'\|_{L_2} = \|\tilde{e}\|_{L_2}$, and

$$\|\tilde{E}''\|_{L_2} = \|\tilde{e}'\|_{L_2} \leq |u^0|_{W^1(L_2(\mathbb{R}))} + |\tilde{v}|_{W^1(L_2(\mathbb{R}))} \leq 2|u^0|_{W^1(L_2(\mathbb{R}))}.$$

Kolmogorov-Landau inequalities in $L^2(\mathbb{R})$ (page 156 in [3]) for the functions $\tilde{E}(x)$, \tilde{E}' , and \tilde{E}'' and the above estimate for $\|\tilde{E}''\|_{L_2}$ give

$$\|\tilde{e}\|_{L_2} = \|\tilde{E}'\|_{L_2} \leq \sqrt{2}\|\tilde{E}\|_{L_2}^{1/2}\|\tilde{E}''\|_{L_2}^{1/2} \leq 2\|\tilde{e}\|_{-1,2}^{1/2}|u^0|_{W^1(L_2(\mathbb{R}))}^{1/2}.$$

Using (41), we derive

$$(42) \quad \|\tilde{e}\|_{L_2} \leq Ch^{5/8}|u^0|_{W^1(L_2(\mathbb{R}))}.$$

By the triangle inequality, we have

$$\|e^N\|_{L_2} \leq \|\tilde{e}\|_{L_2} + \|v^N - \tilde{v}\|_{L_2},$$

and using the estimates (42) and (39) we conclude

$$\|e^N\|_{L_2} \leq Ch^{5/8}|u^0|_{W^1(L_2(\mathbb{R}))}$$

where the constant C depends only on T and α . □

4 Appendix: Proof of Lemma 1

Let us recall the statement of the lemma. We claim, see (11), that

$$C \sum_j (\Delta^2 \delta_j)^2 \geq \sum_j \delta_j^2 - \sum_j (\delta'_j)^2 \geq c \sum_j (\Delta^2 \delta_j)^2,$$

with $C = 3\beta$ and $c = \beta^3/4$, where $\beta = \frac{1}{2}\alpha(1 - \alpha)$.

Let us introduce some notation. Let $y_j = \min(\delta_j, \delta_{j+1})$, $\Delta\delta_j = \delta_j - \delta_{j-1}$, $\Delta y_j = y_j - y_{j-1}$, $\Delta^2\delta_j = \delta_j - 2\delta_{j-1} + \delta_{j-2}$, and $\Delta^2 y_j = y_j - 2y_{j-1} + y_{j-2}$. Recall that $\{\delta_j\}_{-\infty}^{\infty} \in l_2$, and $\delta_j \geq 0$ for all j . It is enough to prove Lemma 1 only for $0 < \alpha < 1$. We construct the new sequence $\{\delta'_j\}$ by using the rule

$$(43) \quad \delta'_j = (1 - \alpha)\delta_j + \alpha\delta_{j-1} - \beta\Delta^2 y_j,$$

for each j . First we assume that $\{\delta_j\}$ has finite support. It is easy to see how to modify the proof in case the support is not finite. Therefore we assume $\delta_j = 0$ for $j \leq 3$ and for $j \geq M - 3$ for some integer M . Then $\delta'_j = 0$ for $j \leq 3$ and $j \geq M - 2$. Thus

$$(44) \quad \|\{\delta_j\}\|_{l_2}^2 - \|\{\delta'_j\}\|_{l_2}^2 = \sum_{j=1}^M \delta_j^2 - \sum_{j=1}^M (\delta'_j)^2.$$

We have

$$\sum_{j=1}^M (\delta'_j)^2 = \sum_{j=1}^M \left(((1-\alpha)\delta_j + \alpha\delta_{j-1})^2 - 2\beta((1-\alpha)\delta_j + \alpha\delta_{j-1})\Delta^2 y_j + \beta^2(\Delta^2 y_j)^2 \right).$$

Note that since $\delta_0 = \delta_1 = 0$ and $\delta_{M-1} = \delta_M = 0$, we have

$$\begin{aligned} & \sum_{j=1}^M \delta_j^2 - ((1-\alpha)\delta_j + \alpha\delta_{j-1})^2 \\ &= \sum_{j=1}^M \left((1 - (1-\alpha)^2)\delta_j^2 - 2\alpha(1-\alpha)\delta_j\delta_{j-1} - \alpha^2 \sum_{j=1}^M \delta_{j-1}^2 \right) \\ &= \sum_{j=1}^M (1 - (1-\alpha)^2 - \alpha^2)\delta_j^2 - 2\alpha(1-\alpha)\delta_j\delta_{j-1} \\ &= 2\beta \sum_{j=1}^M (2\delta_j^2 - 2\delta_j\delta_{j-1}) = 2\beta \sum_{j=1}^M (\delta_j - \delta_{j-1})^2 = 2\beta \sum_{j=1}^M (\Delta\delta_j)^2. \end{aligned}$$

Therefore

$$(45) \quad \sum_{j=1}^M \delta_j^2 - ((1-\alpha)\delta_j + \alpha\delta_{j-1})^2 = 2\beta \sum_{j=1}^M (\Delta\delta_j)^2.$$

Thus we get

$$(46) \quad \sum_{j=1}^M \delta_j^2 - \sum_{j=1}^M (\delta'_j)^2 = 2\beta \sum_{j=1}^M \left((\Delta\delta_j)^2 + ((1-\alpha)\delta_j + \alpha\delta_{j-1})\Delta^2 y_j - \frac{\beta}{2}(\Delta^2 y_j)^2 \right).$$

Now we use $\Delta^2 y_j = \Delta y_j - \Delta y_{j-1}$, $\Delta y_j = 0$, $\delta_j = 0$ for $j \leq 1$, $j \geq M - 1$, and Abel's transform to obtain

$$\sum_{j=1}^M \delta_j \Delta^2 y_j = \sum_{j=1}^M (\delta_j - \delta_{j+1}) \Delta y_j,$$

and

$$\sum_{j=1}^M \delta_{j-1} \Delta^2 y_j = \sum_{j=1}^M (\delta_{j-1} - \delta_j) \Delta y_j.$$

So, (46) becomes

$$\begin{aligned} \sum_{j=1}^M \delta_j^2 - \sum_{j=1}^M (\delta'_j)^2 &= 2\beta \left(\sum_{j=1}^M (\Delta\delta_j)^2 - (1-\alpha) \sum_{j=1}^M \Delta\delta_{j+1}\Delta y_j \right. \\ &\quad \left. - \alpha \sum_{j=1}^M \Delta\delta_j\Delta y_j - \frac{\beta}{2} \sum_{j=1}^M (\Delta^2 y_j)^2 \right) \\ &= 2\beta Q_1 + \beta^2 Q_2, \end{aligned}$$

where

$$Q_1 = \sum_{j=1}^M (\Delta\delta_j)^2 - (1-\alpha) \sum_{j=1}^M \Delta\delta_{j+1}\Delta y_j - \alpha \sum_{j=1}^M \Delta\delta_j\Delta y_j - \beta \sum_{j=1}^M (\Delta^2 \delta_j)^2,$$

and

$$Q_2 = 2 \sum_{j=1}^M (\Delta^2 \delta_j)^2 - \sum_{j=1}^M (\Delta^2 y_j)^2.$$

To finish the proof, it is sufficient prove the following three Lemmas:

Lemma 2. $Q_1 = \frac{1}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j-1} \geq 0} ((1-\alpha)\Delta\delta_{j-1} + \alpha\Delta\delta_j)^2$

$$+ \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0} ((1-\alpha)\Delta\delta_j + \alpha\Delta\delta_{j-1})^2 - \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} \Delta\delta_j \Delta\delta_{j+1}$$

$$+ \left(\frac{1-\alpha}{2} - \beta \right) \sum_{\Delta\delta_{j-1} \geq 0, \Delta\delta_j \geq 0} (\Delta^2 \delta_j)^2 + \left(\frac{\alpha}{2} - \beta \right) \sum_{\Delta\delta_{j-1} < 0, \Delta\delta_j < 0} (\Delta^2 \delta_j)^2 \geq 0.$$

Lemma 3. $Q_2 = \sum_j (((\Delta\delta_{j+1})_- - (\Delta\delta_j)_-) + ((\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+))^2$

$$- 2 \sum_j \Delta\delta_j \Delta\delta_{j-1} (1 - \text{sgn}(\Delta\delta_j \Delta\delta_{j-1})) \geq 0.$$

Here $\text{sgn}(x)$ is the usual sign function defined by $\text{sgn}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$, and

$u_+ = \max(u, 0)$, $u_- = \min(u, 0)$.

Lemma 4. $3\beta \sum_{j=1}^M (\Delta\delta_j)^2 \geq 2\beta Q_1 + \beta^2 Q_2 \geq c \sum_{j=1}^M (\Delta\delta_j)^2.$

The proof of Lemma 2. We consider that \sum_j denotes $\sum_{j=1}^M$. Denote

$$A = \sum_j \Delta\delta_{j+1}\Delta y_j,$$

$$B = \sum_j \Delta\delta_j\Delta y_j.$$

Our aim is to rewrite

$$(47) \quad Q_1 = \sum_j (\Delta\delta_j)^2 - (1 - \alpha)A - \alpha B - \beta \sum_j (\Delta^2\delta_j)^2$$

in the form indicated in Lemma 2.

It is easy to check that

$$(48) \quad \Delta y_j = (\Delta\delta_j)_+ + (\Delta\delta_{j+1})_-.$$

We can transform A as follows:

$$(49) \quad \begin{aligned} A &= \sum_j \Delta\delta_{j+1}((\Delta\delta_j)_+ + (\Delta\delta_{j+1})_-) = \sum_j \Delta\delta_{j+1}(\Delta\delta_j)_+ + \sum_j \Delta\delta_j(\Delta\delta_j)_- \\ &= \sum_{\Delta\delta_j < 0} (\Delta\delta_j)^2 + \sum_{\Delta\delta_j \geq 0} \Delta\delta_j\Delta\delta_{j+1} \\ &= \sum_{\Delta\delta_j < 0} (\Delta\delta_j)^2 + \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} \Delta\delta_j\Delta\delta_{j+1} + D, \end{aligned}$$

where

$$D = \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} \geq 0} \Delta\delta_j\Delta\delta_{j+1}.$$

Further,

$$\begin{aligned} D &= \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} \geq 0} ((\Delta\delta_j)^2 + (\Delta\delta_{j+1})^2 - (\Delta^2\delta_{j+1})^2) \\ &= \frac{1}{2} \sum_{\Delta\delta_{j-1} \geq 0, \Delta\delta_j \geq 0} ((\Delta\delta_{j-1})^2 + (\Delta\delta_j)^2 - (\Delta^2\delta_j)^2) \\ &= \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} \geq 0} (\Delta\delta_j)^2 + \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} \geq 0} (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_{j-1} \geq 0, \Delta\delta_j \geq 0} (\Delta^2\delta_j)^2. \end{aligned}$$

Also, note that

$$(50) \quad \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} \geq 0} (\Delta\delta_j)^2 = \sum_{\Delta\delta_j \geq 0} (\Delta\delta_j)^2 - \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} (\Delta\delta_j)^2$$

and

$$(51) \quad \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} \geq 0} (\Delta\delta_j)^2 = \sum_{\Delta\delta_j \geq 0} (\Delta\delta_j)^2 - \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0} (\Delta\delta_j)^2$$

By (49), (50), and (51) we get

$$(52) \quad A = \sum_j (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0} (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_{j-1} \geq 0, \Delta\delta_j \geq 0} (\Delta^2\delta_j)^2 + \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} \Delta\delta_j \Delta\delta_{j+1}.$$

Transform B in the same way as A :

$$(53) \quad B = \sum_{\Delta\delta_j \geq 0} (\Delta\delta_j)^2 + \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} \Delta\delta_j \Delta\delta_{j+1} + E,$$

where

$$E = \sum_{\Delta\delta_j < 0, \Delta\delta_{j+1} < 0} \Delta\delta_j \Delta\delta_{j+1}.$$

The quantity E can also be rewritten in the same way as D :

$$E = \frac{1}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j+1} < 0} (\Delta\delta_j)^2 + \frac{1}{2} \sum_{\Delta\delta_{j-1} < 0, \Delta\delta_j < 0} (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_{j-1} < 0, \Delta\delta_j < 0} (\Delta^2\delta_j)^2.$$

Combining this equality with (53) we get

$$(54) \quad B = \sum_j (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j+1} \geq 0} (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j-1} \geq 0} (\Delta\delta_j)^2 - \frac{1}{2} \sum_{\Delta\delta_{j-1} < 0, \Delta\delta_j < 0} (\Delta^2\delta_j)^2 + \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} \Delta\delta_j \Delta\delta_{j+1}.$$

By (52) and (54),

$$(55) \quad Q_1 = \sum_j (\Delta\delta_j)^2 - (1-\alpha)A - \alpha B - \beta \sum_j (\Delta^2\delta_j)^2 = \sum_{i=1}^9 S_i,$$

where

$$\begin{aligned} S_1 &= \frac{1-\alpha}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} (\Delta\delta_j)^2, & S_2 &= \frac{1-\alpha}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0} (\Delta\delta_j)^2, \\ S_3 &= \frac{\alpha}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j+1} \geq 0} (\Delta\delta_j)^2, & S_4 &= \frac{\alpha}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j-1} \geq 0} (\Delta\delta_j)^2, \\ S_5 &= - \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j+1} < 0} \Delta\delta_j \Delta\delta_{j+1}, \\ S_6 &= \left(\frac{1-\alpha}{2} - \beta \right) \sum_{\Delta\delta_{j-1} \geq 0, \Delta\delta_j \geq 0} (\Delta^2\delta_j)^2, & S_7 &= \left(\frac{\alpha}{2} - \beta \right) \sum_{\Delta\delta_{j-1} < 0, \Delta\delta_j < 0} (\Delta^2\delta_j)^2, \end{aligned}$$

$$S_8 = -\beta \sum_{\Delta\delta_{j-1} \geq 0, \Delta\delta_j < 0} (\Delta^2\delta_j)^2, \text{ and } S_9 = -\beta \sum_{\Delta\delta_{j-1} < 0, \Delta\delta_j \geq 0} (\Delta^2\delta_j)^2.$$

To get an upper bound for Q_1 note that $|S_j| \leq \frac{1}{2} \sum_j (\Delta^2\delta_j)^2$ for $j = 1, 2, 3, 4, 5$. Therefore, $Q_1 \leq (3 - 2\beta) \sum_j (\Delta^2\delta_j)^2$. (Estimating the relevant quantities more carefully one can replace the constant 3 in the last inequality by $\frac{5}{4}$.)

Also, from the definition of Q_2 , $Q_2 \leq 2 \sum_j (\Delta^2\delta_j)^2$. Combining the upper bounds for Q_1 and Q_2 we get the upper bound in Lemma 1.

Next, note that

$$S_1 + S_4 + S_8 = \sum_{\Delta\delta_j < 0, \Delta\delta_{j-1} \geq 0} \left(\frac{1-\alpha}{2} (\Delta\delta_{j-1})^2 + \frac{\alpha}{2} (\Delta\delta_j)^2 - \beta (\Delta^2\delta_j)^2 \right).$$

Also,

$$\begin{aligned} \frac{1-\alpha}{2} (\Delta\delta_{j-1})^2 + \frac{\alpha}{2} (\Delta\delta_j)^2 - \beta (\Delta^2\delta_j)^2 &= \frac{1-\alpha}{2} (\Delta\delta_{j-1})^2 + \frac{\alpha}{2} (\Delta\delta_j)^2 \\ &\quad - (1-\alpha)\alpha (\Delta\delta_j - \Delta\delta_{j-1})^2 = \frac{1}{2} ((1-\alpha)\Delta\delta_{j-1} + \alpha\Delta\delta_j)^2. \end{aligned}$$

Summing the last inequality over all j with $\Delta\delta_{j-1} \geq 0$, $\Delta\delta_j < 0$, we get

$$(56) \quad S_1 + S_4 + S_8 = \frac{1}{2} \sum_{\Delta\delta_j < 0, \Delta\delta_{j-1} \geq 0} ((1-\alpha)\Delta\delta_{j-1} + \alpha\Delta\delta_j)^2.$$

Similarly,

$$S_2 + S_3 + S_9 = \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0} \left(\frac{1-\alpha}{2} (\Delta\delta_j)^2 + \frac{\alpha}{2} (\Delta\delta_{j-1})^2 - \beta (\Delta^2\delta_j)^2 \right).$$

As before,

$$\frac{1-\alpha}{2} (\Delta\delta_j)^2 + \frac{\alpha}{2} (\Delta\delta_{j-1})^2 - \beta (\Delta^2\delta_j)^2 = \frac{1}{2} ((1-\alpha)\Delta\delta_j + \alpha\Delta\delta_{j-1})^2.$$

Summing the last inequality over all j with $\Delta\delta_j \geq 0$, $\Delta\delta_{j-1} < 0$, we get

$$(57) \quad S_2 + S_3 + S_9 = \frac{1}{2} \sum_{\Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0} ((1-\alpha)\Delta\delta_j + \alpha\Delta\delta_{j-1})^2.$$

This completes the proof of Lemma 2.

The proof of Lemma 3.

By (48) we have $\Delta y_j = (\Delta\delta_j)_+ + (\Delta\delta_{j+1})_-$. We also have $\Delta^2\delta_j = \Delta\delta_j - \Delta\delta_{j-1}$.

Thus, we need to rewrite the quantity

$$Q_2 = 2 \sum_j (\Delta\delta_j - \Delta\delta_{j-1})^2 - \sum_j (((\Delta\delta_{j+1})_- - (\Delta\delta_j)_-) - ((\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+))^2$$

in the form indicated in Lemma 3.

Now, use that for any real numbers x and y , $(x - y)^2 = 2x^2 + 2y^2 - (x + y)^2$. We get

$$\begin{aligned}
& \sum_j (((\Delta\delta_{j+1})_- - (\Delta\delta_j)_-) - ((\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+))^2 \\
&= 2 \left(\sum_j ((\Delta\delta_{j+1})_- - (\Delta\delta_j)_-)^2 + \sum_j ((\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+)^2 \right) \\
&\quad - \sum_j ((\Delta\delta_{j+1})_- - (\Delta\delta_j)_- + (\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+)^2 \\
&= 2 \left(\sum_j ((\Delta\delta_j)_- - (\Delta\delta_{j-1})_-)^2 + \sum_j ((\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+)^2 \right) \\
(58) \quad & - \sum_j ((\Delta\delta_{j+1})_- - (\Delta\delta_j)_- + (\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+)^2
\end{aligned}$$

(in the last equality we have used that the sequence $\{\Delta\delta_j\}$ has finite support.)

Now, we claim that for any real numbers x and y

$$(59) \quad (x - y)^2 + xy(1 - \operatorname{sgn}(xy)) = (x_- - y_-)^2 + (x_+ - y_+)^2.$$

Well, if $x = 0$, both sides of (59) equal y^2 . Similarly, if $y = 0$, both sides of (59) equal x^2 .

If $\operatorname{sgn}(xy) = -1$, then the LHS of (59) is $(x - y)^2 + 2xy$ and the RHS is $x^2 + y^2$.

Finally, if $\operatorname{sgn}(xy) = 1$, then both the LHS and the RHS of (59) equal $(x - y)^2$.

We have shown that (59) holds in all cases.

By using (59) with $x = \Delta\delta_j$, $y = \Delta\delta_{j-1}$, and summing over j we get

$$\begin{aligned}
& \sum_j ((\Delta\delta_j)_- - (\Delta\delta_{j-1})_-)^2 + \sum_j ((\Delta\delta_j)_+ - (\Delta\delta_{j-1})_+)^2 \\
(60) \quad &= \sum_j (\Delta\delta_j - \Delta\delta_{j-1})^2 + \sum_j \Delta\delta_j \Delta\delta_{j-1} (1 - \operatorname{sgn}(\Delta\delta_j \Delta\delta_{j-1})).
\end{aligned}$$

Combining (58) and (60) completes the proof of Lemma 3.

The proof of Lemma 4.

We proved the upper bound of Lemma 4 earlier, so we concentrate on the lower bound.

Denote the five sums appearing in Lemma 2 by $\Sigma_1, \dots, \Sigma_5$, and the two sums appearing in Lemma 3 by Σ_6 , and Σ_7 . Also, let

$$\begin{aligned}
\mathcal{A}_{++} &= \{j : \Delta\delta_j \geq 0, \Delta\delta_{j-1} \geq 0\}, \\
\mathcal{A}_{+-} &= \{j : \Delta\delta_j \geq 0, \Delta\delta_{j-1} < 0\}, \\
\mathcal{A}_{-+} &= \{j : \Delta\delta_j < 0, \Delta\delta_{j-1} \geq 0\}, \text{ and} \\
\mathcal{A}_{--} &= \{j : \Delta\delta_j < 0, \Delta\delta_{j-1} < 0\}.
\end{aligned}$$

Since $\frac{1-\alpha}{2} - \beta = \frac{(1-\alpha)^2}{2} > \frac{\beta^2}{2}$ and $\frac{\alpha}{2} - \beta = \frac{\alpha^2}{2} > \frac{\beta^2}{2}$ we get $\Sigma_4 > \frac{\beta^2}{2} \sum_{j \in \mathcal{A}_{++}} (\Delta^2 \delta_j)^2$, and

$$\Sigma_5 > \frac{\beta^2}{2} \sum_{j \in \mathcal{A}_{--}} (\Delta^2 \delta_j)^2.$$

Next, we need to divide both \mathcal{A}_{+-} and \mathcal{A}_{-+} to a “good” part and a “bad” part.

Define

$$\begin{aligned} \mathcal{A}_{+-}^g &= \{j \in \mathcal{A}_{+-} : |(1-\alpha)\Delta\delta_{j-1} + \alpha\Delta\delta_j| \geq \frac{1}{2} \max((1-\alpha)|\Delta\delta_{j-1}|, \alpha|\Delta\delta_j|)\}, \\ \mathcal{A}_{+-}^b &= \{j \in \mathcal{A}_{+-} : |(1-\alpha)\Delta\delta_{j-1} + \alpha\Delta\delta_j| < \frac{1}{2} \max((1-\alpha)|\Delta\delta_{j-1}|, \alpha|\Delta\delta_j|)\}, \\ \mathcal{A}_{-+}^g &= \{j \in \mathcal{A}_{-+} : |(1-\alpha)\Delta\delta_j + \alpha\Delta\delta_{j-1}| \geq \frac{1}{2} \max((1-\alpha)|\Delta\delta_j|, \alpha|\Delta\delta_{j-1}|)\}, \\ \mathcal{A}_{-+}^b &= \{j \in \mathcal{A}_{-+} : |(1-\alpha)\Delta\delta_j + \alpha\Delta\delta_{j-1}| < \frac{1}{2} \max((1-\alpha)|\Delta\delta_j|, \alpha|\Delta\delta_{j-1}|)\}. \end{aligned}$$

Now, when $j \in \mathcal{A}_{+-}^g$, $|(1-\alpha)\Delta\delta_{j-1} + \alpha\Delta\delta_j| \geq \frac{1}{2} \max((1-\alpha)|\Delta\delta_{j-1}|, \alpha|\Delta\delta_j|) > \beta \max(|\Delta\delta_{j-1}|, |\Delta\delta_j|) \geq \frac{\beta}{2} |\Delta^2 \delta_j|$. Therefore, $\Sigma_2 > \frac{\beta^2}{8} \sum_{j \in \mathcal{A}_{+-}^g} (\Delta^2 \delta_j)^2$. Exactly in the same

way, we get $\Sigma_1 > \frac{\beta^2}{8} \sum_{j \in \mathcal{A}_{-+}^g} (\Delta^2 \delta_j)^2$.

Next use that

$$(61) \quad \text{If } |x+y| < \frac{1}{2} \max(|x|, |y|), \text{ then } \frac{1}{2} < \left| \frac{x}{y} \right| < 2.$$

Indeed, by symmetry it is sufficient to consider the case $|y| \geq |x|$. In this case $|y| - |x| \leq |x+y| < \frac{1}{2} \max(|x|, |y|) = \frac{1}{2}|y|$, and we get $\frac{1}{2}|y| < |x|$, so $1 \leq \left| \frac{x}{y} \right| < 2$, completing the proof of (61).

Next, let $j \in \mathcal{A}_{+-}^b$. By, (61) $\frac{1}{2} < \frac{(1-\alpha)|\Delta\delta_{j-1}|}{\alpha|\Delta\delta_j|} < 2$. So,

$$\frac{(|\Delta\delta_j| + |\Delta\delta_{j-1}|)^2}{|\Delta\delta_j||\Delta\delta_{j-1}|} = \frac{|\Delta\delta_j|}{|\Delta\delta_{j-1}|} + 2 + \frac{|\Delta\delta_{j-1}|}{|\Delta\delta_j|} < \frac{2(1-\alpha)}{\alpha} + 2 + \frac{2\alpha}{1-\alpha} = \frac{1-2\beta}{\beta}.$$

We get

$$(62) \quad (\Delta^2 \delta_j)^2 \leq (|\Delta\delta_j| + |\Delta\delta_{j-1}|)^2 < \frac{1-2\beta}{\beta} |\Delta\delta_j||\Delta\delta_{j-1}|.$$

Exactly in the same way, one can show that (62) holds when $j \in \mathcal{A}_{-+}^b$. Also, note that $\mathcal{A}_{+-}^b \cap \mathcal{A}_{-+}^b = \emptyset$ and if $j \in \mathcal{A}_{+-}^b \cup \mathcal{A}_{-+}^b$, then $\text{sgn}(\Delta\delta_j \Delta\delta_{j-1}) = -1$. We get,

$$\Sigma_7 > 4\beta \left(\sum_{j \in \mathcal{A}_{+-}^b} (\Delta^2 \delta_j)^2 + \sum_{j \in \mathcal{A}_{-+}^b} (\Delta^2 \delta_j)^2 \right).$$

Combining the lower bounds for $\Sigma_4, \Sigma_5, \Sigma_2, \Sigma_1$, and Σ_7 we get the lower bound in Lemma 4 (note that all Σ 's are non-negative). This completes the proof of Lemma 4 and of Lemma 1.

References

- [1] Y. Brenier and S. Osher, The one-sided Lipschitz condition for convex scalar conservation laws, *SIAM J. Numer. Anal.*, 25: 8–23, 1988.

- [2] P. Brenner, V. Thomée and L. B. Wahlbin, Besov spaces and applications to difference methods for initial value problems, (A. Dold and B. Eckmann, eds.) *Lecture Notes in Math.*, vol. 434, Springer-Verlag, Berlin and New York, 1975.
- [3] R. A. DeVore and G. G. Lorentz, *Constructive Approximation*, Springer-Verlag, Berlin, 1993.
- [4] A. Harten and S. Osher, Uniformly high order accurate non-oscillatory schemes, I, *J. Appl. Num. Math.*, volume 71, 2: 279–309, 1987.
- [5] A. Harten, B. Enquist, S. Osher and S.R. Chakravarthy, Uniformly high order accurate essentially non-oscillatory schemes, III, *J. Comp. Phys.*, volume 71, 2: 231–303, 1987.
- [6] G.-S. Jiang, D. Levi, C.-T. Lin, S. Osher and E. Tadmor High-resolution non-oscillatory central schemes with nonstaggered grids for hyperbolic conservation laws. *SIAM J. Numer. Anal.*, 35, 6: 2147–2169, 1998.
- [7] G.-S. Jiang and E. Tadmor Nonoscillatory central schemes for hyperbolic conservation laws. *SIAM J. Sci. Comput.*, 19, 6: 1892–1917, 1998.
- [8] Yu. V. Kryakin On the theorem of H. Whitney in spaces L^p , $1 \leq p \leq \infty$. *Mathematica Balkanica*, New series, Vol. 4, Fasc. 3: 258–270, 1990.
- [9] P. Lax and B. Wendroff, Systems of conservation laws, *Comm. Pure Appl. Math.* 13: 217–237, 1960.
- [10] S. Konyagin, B. Popov and O. Trifonov, On Convergence of Minmod-Type Schemes, submitted to SIAM Numerical Analysis,
- [11] S.N. Kruzhkov, First order quasi-linear equations in several independent variables, *Math. USSR Sbornik*, Vol. 10, #2: 217–243, 1970.
- [12] A. Kurganov and E. Tadmor, New high-resolution central schemes for nonlinear conservation laws and convection-diffusion equations, *J. Comp. Phys.*, volume 160: 241–282, 2000.
- [13] H. Nessyahu and E. Tadmor, Non-oscillatory central differencing for hyperbolic conservation laws, *J. Comp. Phys.*, volume 87, 2: 408–463, 1990.
- [14] T. Tang and Z.-H. Teng, The sharpness of Kuznetsov’s $O(\sqrt{\Delta x})$ L^1 -error estimate for monotone difference scheme, *Math. Comp.*, 64: 581–589, 1995.
- [15] C.-W. Shu, Numerical experiments on the accuracy of ENO and modified ENO schemes, *J. Comp. Phys.*, 5: 127–149, 1990.