



INDUSTRIAL
MATHEMATICS
INSTITUTE

2006:10

Littlewood-Paley theorem for
Schrodinger operators

Shijun Zheng

IMI

Preprint Series

Department of Mathematics
University of South Carolina

LITTLEWOOD-PALEY THEOREM FOR SCHRÖDINGER OPERATORS

SHIJUN ZHENG

ABSTRACT. Let H be a Schrödinger operator on \mathbb{R}^n . Under a polynomial decay condition for the kernel of its spectral operator, we show that the Besov spaces and Triebel-Lizorkin spaces associated with H are well defined. We further give a Littlewood-Paley characterization of L_p spaces in terms of dyadic functions of H . This generalizes and strengthens the previous result when the heat kernel of H satisfies certain upper Gaussian bound.

1. INTRODUCTION AND MAIN RESULTS

Recently the theory of function spaces associated with Schrödinger operators have been drawing attention in the area of harmonic analysis and PDEs [12, 2, 1, 14, 16, 9, 6, 8, 7, 5]. In [9, 6, 1, 14] it is proved that the Besov and Triebel-Lizorkin spaces associated with a Schrödinger operator are well defined, in some particular cases. In this note we aim to extend the result for general Schrödinger operators on \mathbb{R}^n . Furthermore we are interested in obtaining a Littlewood-Paley decomposition for the L_p spaces as well as Sobolev spaces using dyadic functions of H .

Let $H = -\Delta + V$ be a Schrödinger operator that is selfadjoint in $L_2(\mathbb{R}^n)$ with a real-valued potential function V . Then for a Borel measurable function ϕ , one can define the spectral operator $\phi(H)$ by functional calculus $\phi(H) = \int_{-\infty}^{\infty} \phi(\lambda) dE_\lambda$, where dE_λ is the spectral measure of H . The kernel of $\phi(H)$ is denoted $\phi(H)(x, y)$.

Let $\{\varphi_j\}_{j \in \mathbb{Z}} \subset C_0^\infty(\mathbb{R})$ be a smooth dyadic system satisfying the conditions (i) $\text{supp } \varphi_j \subset \{x : 2^{j-2} \leq |x| \leq 2^j\}$

$$(ii) |\varphi_j^{(k)}(x)| \leq c_k 2^{-kj}, \quad \forall j \in \mathbb{Z}, k \in \mathbb{N}_0 = \{0\} \cup \mathbb{N}$$

Date: July 26, 2006.

2000 Mathematics Subject Classification. Primary: 42B25; Secondary: 35P25.

Key words and phrases. functional calculus, Schrödinger operator, Littlewood-Paley theory.

This work is supported by DARPA grant HM1582-05-2-0001. The author gratefully thanks the hospitality and support of Department of Mathematics, University of South Carolina, during his visiting at the Industrial Mathematics Institute.

(iii)

$$\sum_{j=-\infty}^{\infty} \varphi_j(x) \approx c > 0, \quad \forall x \neq 0.$$

Let $0 < p < \infty$, $0 < q \leq \infty$ and $\alpha \in \mathbb{R}$. The homogenous *Triebel-Lizorkin space* $\dot{F}_p^{\alpha,q}(H)$ is defined as the completion of the Schwartz class $\mathcal{S}(\mathbb{R}^n)$ with the quasi-norm

$$\|f\|_{\dot{F}_p^{\alpha,q}(H)} = \left\| \left(\sum_{j=-\infty}^{\infty} 2^{j\alpha q} |\varphi_j(H)f(\cdot)|^q \right)^{1/q} \right\|_p.$$

Similarly, if $0 < p \leq \infty$, $0 < q \leq \infty$, the homogeneous *Besov space* $\dot{B}_p^{\alpha,q}(H)$ is defined by the quasi-norm

$$\|f\|_{\dot{B}_p^{\alpha,q}(H)} = \left(\sum_{j=-\infty}^{\infty} 2^{j\alpha q} \|\varphi_j(H)f\|_p^q \right)^{1/q}.$$

Throughout this note we assume H satisfies the following:

Assumption 1.1. *Let $\phi_j \in C_0^\infty(\mathbb{R})$ be as in condition (i), (ii). Then for every $N \in \mathbb{N}_0$ there exists a constant $c_N > 0$ such that for all $j \in \mathbb{Z}$*

$$(1) \quad |\phi_j(H)(x, y)| \leq c_N \frac{2^{nj/2}}{(1 + 2^{j/2}|x - y|)^N}$$

$$(2) \quad |\nabla_x \phi_j(H)(x, y)| \leq c_N \frac{2^{(n+1)j/2}}{(1 + 2^{j/2}|x - y|)^N}.$$

This is the case when H is the Hermite operator $-\Delta + |x|^2$, or more generally, whenever V is nonnegative and H satisfies the upper Gaussian bound for the heat kernel and its derivative (see Proposition 3.3). However, when the potential V is negative, such a heat kernel estimate is *not* available. Therefore it is necessary to consider a more general condition as given in Assumption 1.1.

Define the Peetre maximal function for H as: for $j \in \mathbb{Z}$, $s > 0$

$$\varphi_{j,s}^* f(x) = \sup_{t \in \mathbb{R}^n} \frac{|\varphi_j(H)f(t)|}{(1 + 2^{j/2}|x - t|)^s},$$

and

$$\varphi_{j,s}^{**} f(x) = \sup_{t \in \mathbb{R}^n} \frac{|(\nabla_t \varphi_j(H)f)(t)|}{(1 + 2^{j/2}|x - t|)^s}.$$

The following theorem is a maximal characterization of the *homogeneous spaces*. By $\|\cdot\|_A \approx \|\cdot\|_B$ we mean equivalent norms.

Theorem 1.2. *Suppose H satisfies Assumption 1.1.*

a) *If $0 < p \leq \infty$, $0 < q \leq \infty$, $\alpha \in \mathbb{R}$ and $s > n/p$, then*

$$\|f\|_{\dot{B}_p^{\alpha,q}(H)} \approx \|\{2^{j\alpha}\varphi_{j,s}^*(H)f\}\|_{\ell^q(L_p)}.$$

b) *If $0 < p < \infty$, $0 < q \leq \infty$, $\alpha \in \mathbb{R}$ and $s > n/\min(p, q)$, then*

$$\|f\|_{\dot{F}_p^{\alpha,q}(H)} \approx \|\{2^{j\alpha}\varphi_{j,s}^*(H)f\}\|_{L_p(\ell^q)}.$$

It is well-known that such a characterization implies that any two dyadic systems satisfying (i), (ii), (iii) give rise to equivalent norms on $\dot{F}_p^{\alpha,q}(H)$ and $\dot{B}_p^{\alpha,q}(H)$. The analogous result also holds for the inhomogeneous spaces $F_p^{\alpha,q}(H)$, $B_p^{\alpha,q}(H)$. However, the homogeneous spaces, which cover both high and low energy portion of H , are essential and more useful in proving Strichartz inequality for wave equations [16, 13]. This is one reason of our motivation.

Following the same idea in [14], using Calderón-Zygmund decomposition and Assumption 1.1 we show that $L_p(\mathbb{R}^n) = F_p^{0,2}(H)$ if $1 < p < \infty$. We thus obtain the Littlewood-Paley theorem for L_p spaces.

Theorem 1.3. *Suppose H satisfies Assumption 1.1. If $1 < p < \infty$, then*

$$\|f\|_{L_p(\mathbb{R}^n)} \approx \left\| \left(\sum_{j=-\infty}^{\infty} |\varphi_j(H)f(\cdot)|^2 \right)^{1/2} \right\|_{L_p(\mathbb{R}^n)}.$$

Under additional condition on V , e.g., $|\partial_x^k V(x)| \leq c_k$, $|k| \leq 2m_0 - 2$ for some $m_0 \in \mathbb{N}$, we can characterize the Sobolev spaces $H_p^{2s}(\mathbb{R}^n) = F_p^{s,2}(H)$, $1 < p < \infty$, $|s| \leq m_0$ with equivalent norms

$$\|f\|_{H_p^{2s}(\mathbb{R}^n)} \approx \left\| \left(\sum_{j=-\infty}^{\infty} 2^{2js} |\varphi_j(H)f(\cdot)|^2 \right)^{1/2} \right\|_{L_p(\mathbb{R}^n)}.$$

2. PROOFS OF THEOREM 1.2 AND THEOREM 1.3

The proof of Theorem 1.2 is standard and follows from Bernstein type inequality (Lemma 2.1) and Peetre type maximal inequality (Lemma 2.2) for maximal functions.

Lemma 2.1. *For $s > 0$, there exists a constant $c_{n,s} > 0$ such that for all $j \in \mathbb{Z}$*

$$\varphi_{j,s}^{**} f(x) \leq c_{n,s} 2^{j/2} \varphi_{j,s}^* f(x), \quad \forall f \in \mathcal{S}(\mathbb{R}^n).$$

Similar to [20, 14] Lemma 2.1 can be easily proved using (2) with $N > n + s$ and the identity

$$\varphi_j(H)f(x) = \psi_j(H)\varphi_j(H)f,$$

where $\psi_j(x) = \psi(2^{-j}x)$ with $\psi \in C_0^\infty$, $\text{supp } \psi \subset \{\frac{1}{5} \leq |x| \leq \frac{5}{4}\}$ and $\psi(x) = 1$ on $\{\frac{1}{4} \leq |x| \leq 1\}$.

Let M denote the Hardy-Littlewood maximal function

$$(3) \quad Mf(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy$$

where the supreme is taken over all balls B in \mathbb{R}^n centered at x .

Lemma 2.2. *Let $0 < r < \infty$ and $s = n/r$. Then for all $j \in \mathbb{Z}$*

$$(4) \quad \varphi_{j,s}^* f(x) \leq c_{n,r} [M(|\varphi_j(H)f|^r)]^{1/r}(x), \quad \forall f \in \mathcal{S}(\mathbb{R}^n).$$

Proof. Let $g(x) \in C^1(\mathbb{R}^n)$. As in [20, 1], the mean value theorem gives for $z_0 \in \mathbb{R}^n$, $\delta > 0$

$$|g(z_0)| \leq \delta \sup_{|z-z_0| \leq \delta} |\nabla g(z)| + c_{n,r} \delta^{-n/r} \left(\int_{|z-z_0| \leq \delta} |g|^r dz \right)^{1/r}.$$

Put $g(z) = \varphi_j(H)f(x-z)$ to get

$$\begin{aligned} \frac{|\varphi_j(H)f(x-z)|}{(1+2^{j/2}|z|)^{n/r}} &\leq \delta \sup_{|u-z| \leq \delta} \frac{(1+2^{j/2}|u|)^{n/r} |\nabla(\varphi_j(H)f)(x-u)|}{(1+2^{j/2}|z|)^{n/r} (1+2^{j/2}|u|)^{n/r}} \\ &+ c_{n,r} \delta^{-n/r} (1+2^{j/2}|z|)^{-n/r} \left(\int_{|u-z| \leq \delta} |\varphi_j(H)f(x-u)|^r du \right)^{1/r} \\ &\leq \delta (1+2^{j/2}\delta)^{n/r} \varphi_{j,s}^{**} f(x) + c_{n,r} \delta^{-n/r} \left(\frac{|z|+\delta}{1+2^{j/2}|z|} \right)^{n/r} [M(|\varphi_j(H)f|^r)(x)]^{1/r} \\ &\leq c_{n,r} \delta (1+2^{j/2}\delta)^{n/r} 2^{j/2} \varphi_{j,s}^* f(x) + c_{n,r} \delta^{-n/r} \left(\frac{|z|+\delta}{1+2^{j/2}|z|} \right)^{n/r} [M(|\varphi_j(H)f|^r)]^{1/r}(x) \\ &\leq c_{n,r} \epsilon (1+\epsilon)^{n/r} \varphi_{j,s}^* f(x) + c_{n,r} (1+\epsilon^{-1})^{n/r} [M(|\varphi_j(H)f|^r)]^{1/r}(x), \end{aligned}$$

by setting $\delta = 2^{-j/2}\epsilon$, $\epsilon > 0$ and using Lemma 2.1. Finally, taking $\epsilon > 0$ sufficiently small establishes (4). \square

Now Theorem 1.2 is a consequence of Lemma 2.2 and the following well-known lemma on Hardy-Littlewood maximal function by a standard argument; see [20] or [9, 14] for some simple details.

Lemma 2.3. *a) If $1 < p \leq \infty$, then*

$$(5) \quad \|Mf\|_{L_p(\mathbb{R}^n)} \leq C_p \|f\|_{L_p(\mathbb{R}^n)}.$$

b) If $1 < p < \infty$, $1 < q \leq \infty$, then

$$(6) \quad \left\| \left(\sum_j |Mf_j|^q \right)^{1/q} \right\|_{L_p(\mathbb{R}^n)} \leq C_{p,q} \left\| \left(\sum_j |f_j|^q \right)^{1/q} \right\|_{L_p(\mathbb{R}^n)}.$$

2.4. Proof of Theorem 1.3. From the proof of the identification of $F_p^{0,2}(H)$ spaces [14, Theorem 5.1] we observe that the estimates in (1), (2) imply

$$(7) \quad \|f\|_{F_p^{0,2}(H)} \approx \|f\|_{L_p}, \quad 1 < p < \infty$$

by applying $L_p(\ell^2)$ -valued Calderón-Zygmund decomposition. On the other hand, Theorem 1.2 suggests that

$$(8) \quad \|f\|_{F_p^{\alpha,q}(H)} \approx \|\{2^{j\alpha}\varphi_j(H)f\}\|_{L_p(\ell^q)}$$

whenever $\{\varphi_j\}_{j \in \mathbb{Z}}$ is a dyadic system satisfying (i), (ii), (iii).

Combining (7) and (8) with $\alpha = 0$, $q = 2$ proves Theorem 1.3. \square

Remark 2.5. For $p = 1$, Dziubański and Zienkiewicz [7] recently obtained a characterization of Hardy space associated with H and showed that if a compactly supported positive potential V is in $L^{n/2+\epsilon}$, $n \geq 3$, then

$$\|f\|_{\mathcal{H}^1} \approx \|wf\|_{H^1(\mathbb{R}^d)},$$

where $\mathcal{H}^1 = \{f \in L^1 : \sup_{t>0} |e^{-tH}f(\cdot)| \in L^1\}$ and the weight w is defined by $w(x) = \lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} e^{-tH}(x, y) dy$. It would be very interesting to see whether one can give a Littlewood-Paley characterization of \mathcal{H}^1 in the sense of Theorem 1.3.

3. POTENTIALS SATISFYING UPPER GAUSSIAN BOUND

In this section we show that Assumption 1.1 is verified when H satisfies the upper Gaussian bound (10) for its heat kernel. We begin with a weighted L^1 inequality, which is an easy consequence of [11, Lemma 8] by a scaling argument.

Lemma 3.1. (Hebisch) Suppose $V \geq 0$ and e^{-tH} satisfies

$$(9) \quad 0 \leq e^{-tH}(x, y) \leq c_n t^{-n/2} e^{-c|x-y|^2/t}, \quad \forall t > 0.$$

If $s > (n+1)/2 + \beta$, $\beta \geq 0$ and $\text{supp } g \subset [-10, 10]$, then

$$\sup_{j \in \mathbb{Z}, y \in \mathbb{R}^n} \|g(2^{-j}H)(\cdot, y) \langle 2^{j/2}(\cdot - y) \rangle^\beta\|_{L^1(\mathbb{R}^n)} \leq c_n \|g\|_{H^s(\mathbb{R})},$$

where $\langle x \rangle := 1 + |x|$ and $\|\cdot\|_{H^s}$ denotes the usual Sobolev norm.

Remark 3.2. It is known that (9) holds whenever $V \geq 0$ is locally integrable.

Proposition 3.3. *Let $\alpha = 0, 1$. Suppose $V \geq 0$ and e^{-tH} satisfies the upper Gaussian bound*

$$(10) \quad |\nabla_x^\alpha e^{-tH}(x, y)| \leq c_n t^{-(n+\alpha)/2} e^{-c|x-y|^2/t}, \quad \forall t > 0.$$

If $\{\varphi_j\}_{j \in \mathbb{Z}}$ is a dyadic system satisfying (i), (ii), then for each $N \geq 0$

$$|\nabla_x^\alpha \varphi_j(H)(x, y)| \leq c_N 2^{j(n+\alpha)/2} (1 + 2^{j/2}|x-y|)^{-N}, \quad \forall j.$$

Proof. Write

$$\nabla_x^\alpha \varphi_j(H)(x, y) = \int_z \nabla_x^\alpha e^{-tH}(x, z) (e^{tH} \varphi_j(H))(z, y) dz.$$

By (10) we have

$$\begin{aligned} & |\nabla_x^\alpha \varphi_j(H)(x, y)| \\ & \leq c_n t^{-(n+\alpha)/2} \int e^{-c|x-z|^2/t} \langle (x-z)/\sqrt{t} \rangle^N \langle (x-z)/\sqrt{t} \rangle^{-N} \langle (z-y)/\sqrt{t} \rangle^{-N} \\ & \quad \cdot \langle (z-y)/\sqrt{t} \rangle^N |(e^{tH} \varphi_j(H))(z, y)| dz \\ & \leq c_n t^{-(n+\alpha)/2} \langle (x-y)/\sqrt{t} \rangle^{-N} \int \langle (z-y)/\sqrt{t} \rangle^N |(e^{tH} \varphi_j(H))(z, y)| dz. \end{aligned}$$

Setting $t = t_j := 2^{-j}$, we see that $g_j(x) := e^{t_j x} \varphi_j(x)$ also satisfies conditions (i), (ii). Writing $g_j(x) = g_0(2^{-j}x)$, then $\text{supp } g_0 \subset \{\frac{1}{4} \leq |x| \leq 1\}$ and

$$\|g_0\|_{H^N(\mathbb{R})} \leq \|g_j(2^j x)\|_{C^N(\mathbb{R})} \leq c_N.$$

Thus an application of Lemma 3.1 with $g = g_0$, $\beta = N$ proves the proposition. \square

3.4. Hermite operator $H = -\Delta + |x|^2$. To verify Assumption 1.1 it is sufficient to show e^{-tH} satisfies the upper Gaussian bound in (10), according to Proposition 3.3.

For $k \in \mathbb{N}_0$, let h_k be the k^{th} Hermite function with $\|h_k\|_{L_2(\mathbb{R})} = 1$ such that

$$\left(-\frac{d^2}{dx^2} + x^2\right)h_k = (2k+1)h_k.$$

Then $\{h_k\}_0^\infty$ forms a complete orthonormal system (ONS) in $L_2(\mathbb{R})$. In $L_2(\mathbb{R}^n)$, the ONS is given by $\Phi_k(x) := h_{k_1} \otimes \cdots \otimes h_{k_n}$, $k = (k_1, \dots, k_n) \in \mathbb{N}_0^n$.

By Mehler's formula [18, Ch.4] or [19], the heat kernel has the expression

$$\begin{aligned} e^{-tH}(x, y) &= \sum_{k \in \mathbb{N}_0^n} e^{-t(n+2|k|)} \Phi_k(x) \Phi_k(y) \\ &= \frac{1}{(2\pi \sinh(2t))^{n/2}} e^{-\frac{1}{2} \coth(2t)(|x|^2+|y|^2) + \operatorname{cosech}(2t)x \cdot y} \end{aligned}$$

for all $t > 0$, $x, y \in \mathbb{R}^n$.

It is easy to calculate to find that there exist constants $c, c' > 0$, $0 < c_0, c_1, c'_0, c'_1 < 1$ and $t_0 > 1$ such that

$$\begin{aligned} p_{\frac{t}{2}}(x, y) &\leq c \begin{cases} t^{-n/2} e^{-c_0|x-y|^2/t} & t \leq t_0 \\ e^{-nt/2} e^{-c_1|x-y|^2} & t > t_0 \end{cases} \\ |\nabla_x p_{\frac{t}{2}}(x, y)| &\leq c' \begin{cases} t^{-(n+1)/2} e^{-c'_0|x-y|^2/t} & t \leq t_0 \\ e^{-nt/2} e^{-c'_1|x-y|^2} & t > t_0, \end{cases} \end{aligned}$$

where $p_t(x, y) := e^{-tH}(x, y)$. Hence (10) holds.

Remark 3.5. *For the Hermite operator, the decay estimates similar to (1), (2) were previously obtained in [10] in one dimension and [6] in n -dimension. The latter used Heisenberg group method. Proposition 3.3 shows that using heat kernel estimate we can obtain a simpler proof.*

Remark 3.6. *When V is negative, the heat kernel estimate (9) is not available, especially in low dimensions $n = 1, 2$. but Assumption 1.1 still holds in the high energy case ($j \geq 0$) for certain short range potentials. A special example is the one dimensional Pöschl-Teller model $V(x) = -\nu(\nu + 1) \operatorname{sech}^2 x$, $\nu \in \mathbb{N}$, cf. [14]. We will discuss the problem in more detail in [23] where V is assumed to have only polynomial decay at infinity.*

REFERENCES

- [1] J. Benedetto, S. Zheng, Besov spaces for the Schrödinger operator with barrier potential (submitted). <http://lanl.arXiv.org/math.CA/0411348>, (2005).
- [2] P. D'Ancona, V. Pierfelice, On the wave equation with a large rough potential, *J. Funct. Anal.* **227** (2005), no. 1, 30-77.
- [3] E. Davies, *Heat Kernels and Spectral Theory*, Cambridge University Press, Cambridge, 1989.
- [4] ———, Pointwise bounds on the space and time derivatives of heat kernels, *J. Operator Theory* **21** (1989), 367-378.
- [5] X. Duong, L. Yan, Duality of Hardy and BMO spaces associated with operators with heat kernel bounds, *J. Amer. Math. Soc.* **18** (2005), 943-973.
- [6] J. Dziubański, Triebel-Lizorkin spaces associated with Laguerre and Hermite expansions, *Proc. Amer. Math. Soc.* **125** (1997), 3547-3554.

- [7] J. Dziubański, J. Zienkiewicz, Hardy spaces H^1 for Schrödinger operators with compactly supported potentials, *Ann. Mat. Pura Appl.* (4) **184** (2005), no. 3, 315–326.
- [8] J. Dziubański and J. Zienkiewicz, H^p spaces for Schrödinger operators, in: *Fourier Analysis and Related Topics*, Banach Center Publ. **56**, Inst. Math., Polish Acad. Sci., 2002, 45-53.
- [9] J. Epperson, Triebel-Lizorkin spaces for Hermite expansions, *Studia Math.* **114** (1995), 87-103.
- [10] ———, Hermite and Laguerre wave packet expansions, *Studia Math.* **126** (1997), no. 3, 199–217.
- [11] W. Hebisch, A multiplier theorem for Schrödinger operators, *Colloq. Math.* **60/61** (1990), no. 2, 659-664.
- [12] A. Jensen, S. Nakamura, Mapping properties of functions of Schrödinger operators between L^p spaces and Besov spaces, in *Spectral and Scattering Theory and Applications*, Advanced Studies in Pure Math. **23**, 1994.
- [13] M. Keel, T. Tao, Endpoint Strichartz estimates, *Amer. J. Math.* **120** (1998), 955-980.
- [14] G. Ólafsson, S. Zheng, Function spaces associated with Schrödinger operators: the Pöschl-Teller potential (submitted).
- [15] I. Rodnianski, T. Tao, Long-time decay estimates for the Schrödinger equation on manifolds. *Preprint*.
- [16] W. Schlag, A remark on Littlewood-Paley theory for the distorted Fourier transform, <http://lanl.arXiv.org/math.AP/0508577>, (2005).
- [17] B. Simon, Schrödinger semigroups, *Bull. Amer. Math. Soc.* **7** (1982) no.3, 447-526.
- [18] S. Thangavelu, *Lectures on Hermite and Laguerre expansions*, Princeton Univ. Press, 1993.
- [19] ———, Hermite and Laguerre semigroups, some recent developments. *Preprint*.
- [20] H. Triebel, *Theory of Function Spaces*, Birkhäuser Verlag, 1983.
- [21] —, *Theory of Function Spaces II*, Monographs Math. **84**, Birkhäuser, Basel, 1992.
- [22] S. Zheng, A representation formula related to Schrödinger operators, *Anal. Theo. Appl.* **20** (2004), no.3., 294-296.
- [23] ———, Spectral multipliers, function spaces and dispersive estimates for Schrödinger operators. *Preprint*.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SC 29208

E-mail address: shijun@math.sc.edu

URL: <http://www.math.sc.edu/~shijun>